

CARDIAC EMERGENCIES AND HEART FAILURE

Prevention and Treatment

BY

ARTHUR M MASTER M D

Cardiologist Mount Sinai Hospital New York N Y

MARVIN MOSER M D

1st Lt USAF (MC) Walter Reed Army Hospital Washington D C

Former Fellow Cardiology Mount Sinai Hospital New York N Y

AND

HARRY L JAFFE M D

Adjunct Physician Cardiology Mount Sinai Hospital New York N Y

Illustrated



LEA & FEBIGER

Philadelphia

1952

LEA & FEBIGER

1952

Printed in the United States of America

Preface

THE prompt diagnosis of cardiovascular emergencies and their proper treatment are among the most common problems confronting the general practitioner. This monograph has been written in the hope that it will be of value to him as he encounters these acute conditions. It is not intended to be a book on Cardiology, Radiology, or Electrocardiography; consequently it does not cover these fields as thoroughly as do the standard textbooks. Physiology and pathology are discussed when a fuller understanding of them is considered necessary for the intelligent handling of cardiac patients. The drugs that have been found valuable are stressed; other drugs in current use are discussed. Case histories are presented to illustrate specific conditions and the use of the newer drugs. Throughout the book the importance of preventing cardiac emergencies by adequate interval therapy is emphasized.

We have included sections on surgical cardiac emergencies, on cardiac resuscitation and on certain acute conditions that are not commonly encountered, i.e., dissecting aneu-

rysm hypertensive crises secondary to pheochromocytoma and the acute episode resulting from a "ball valve" thrombus in the hope that when these emergencies are seen they may be recognized and treated properly

The section on congestive heart failure and its complications is brief but we have included all of the information that we believe is necessary for the treatment of this condition

We have surveyed the literature but have drawn freely from our own cases and experience as well as from material previously published by one of us (A. M. M.). Portions of this monograph have been presented in several broadcasts sponsored by the New York Academy of Medicine over Station WNYC (FM)

We wish to thank Dr. Samuel Kahn and Dr. Phillip Samet for editorial assistance. We are indebted to members of the cardiographic department of the Mount Sinai Hospital for valuable suggestions

We hope that the great number of physicians who are called upon to diagnose and treat the diseases that we discuss will benefit from this work

Contents

CHAPTER	PAGE
1 Arrhythmias	11
Paroxysmal Tachycardias	11
Auricular Fibrillation	14
Auricular Flutter	25
Auricular and Nodal (Supraventricular)	
Tachycardia	28
Ventricular Tachycardia	34
Ventricular Fibrillation	39
Complete Heart Block	42
Other Arrhythmias	43
Sinus Tachycardia	43
Sinus Bradycardia	43
Premature Beats	44
2 Acute Pulmonary Edema and Congestive Heart Failure	45
Pathogenesis of Cardiac Failure	45

Paroxysmal Nocturnal Dyspnea	48
Acute Pulmonary Edema	49
Therapy of Congestive Heart Failure	52
3 Angina Pectoris Coronary Insufficiency and Coronary Occlusion	61
Differential Diagnosis	61
Treatment	61
Angina Pectoris	62
Acute Coronary Insufficiency	65
Acute Coronary Occlusion	70
4 Syncope	91
5 Rheumatic Heart Disease	99
6 Hypertensive Encephalopathy and Crises	106
7 Dissecting Aneurysm	108
8 Traumatic Heart Disease	111
Penetrating Chest Injuries	111
Nonpenetrating Chest Injuries	113
9 Surgical Emergencies in Cardiovascular Dis- ease	116
Surgery in the Cardiac Patient	116
Cardiac Arrest	119

List of Illustrations

FIGURE		PAGE
1	<i>Effect of Potassium in Digitalis Toxicity</i>	23
2	Auricular Flutter <i>Effect of Intramuscular Quinidine</i>	26
3	2 1 Auricular Tachycardia	29
4	Nodal Tachycardia <i>Effect of Pronestyl</i>	33
5	Ventricular Tachycardia <i>Effect of Pronestyl</i>	36
6	Complete A V Block and Ventricular Fibrillation	40
7	Acute Coronary Insufficiency Following Ligation	66
8	Acute Coronary Insufficiency Caused by Hemorrhage	68
9	Electrocardiographic Changes of Coronary Insufficiency During the Premonitory Phase of <i>Coronary Occlusion with Anterior Infarction</i>	71
10	Acute Coronary Occlusion with Posterior Infarction	72
11	Pulmonary Embolism with Coronary Insufficiency	82
12	<i>Effect of Carotid Sinus Pressure</i>	94
13	Acute Pericarditis	102

CARDIAC EMERGENCIES AND HEART FAILURE

Introduction

DUE to the tremendous advances in the prevention and treatment of many diseases, and the increased span of life, cardiovascular disease now greatly outnumbers all others, and causes more than half the deaths after the age of fifty. Acute heart conditions require emergency treatment more than do those of any other organ making the term 'acute heart' even more appropriate than the expression 'acute abdomen'. The prompt treatment of acute cardiac disease by the application of well established rules and the utilization of recent advances in the field of cardiology, will certainly save many lives and hasten recovery in others. Since treatment must be immediate, it is incumbent upon the physician to know how to diagnose and treat cardiac emergencies, and to have the essential drugs readily available. This requires the careful selection and constant replenishment of medical supplies for it is very disheartening to make the correct diagnosis of a cardiac emergency, and not have the suitable medication available. The following materials should always be at hand:

1. Aminophyllin 2 cc ampules = 0.48 gm for i m, 10 cc ampules = 24 gm for i v use or 5 gm suppositories
2. Atropin Sulfate 75 mg (gr 1/100) tablets for intravenous or oral use
3. Caffeine Sodium Benzoate 2 cc ampules (5 gm) for i m use

- 4 Cedilanid 4 cc ampules 8 mg (Lanatocide C)
- 5 Demerol 100 mg tablets
- 6 Digoxin 25 mg tablets, 1 cc = 5 mg ampules (must be diluted in 10 cc saline for i v use)
- 7 Dilaudid hypo tablets 2 mg (gr 1/32) or ampules 1 cc = 2 mg (gr 1/32)
- 8 Epinephrine 1 1000—(1 cc)
- 9 Magnesium Sulfate 10 cc of 20 per cent solution ampules
- 10 Mercurial Diuretic (Mercurhydrin or Thiomerin) 1 cc or 2 cc
- 11 Morphine Sulfate 8 mg (1/8 gr) or 16 mg (1/4 gr) tablets for i m or i v use
- 12 Needles for phlebotomy
- 13 Neosynephrin (ampules for i v use) 1 cc = 10 mg
- 14 Nitroglycerin 30 mg (gr 1/200) or 40 mg (gr 1/150) hypodermic tablets
- 15 Potassium Chloride 20 cc ampules = 3.7 gm tablets 3 gm (5 gr)
- 16 Pronestyl (Procaine Amide) 250 mg capsules 10 cc ampules containing 100 mg per cc
- 17 Quinidine 2 gm (3 gr) and 3 gm (5 gr) tablets 2 gm (1 cc) ampules for i m use
- 18 Sodium Amytal 0.6 gm (gr 1) and 2 gm (gr 3) capsules for oral use 5 gm of powder in ampules for parenteral use
- 19 Sterile water for dilution of tablets
- 20 Strophanthin (K Strophanthin) 1 cc = 65 mg or Ouabain (G Strophanthin) 1 cc = 25 mg
- 21 Syrup of Ipecac
- 22 Tourniquets

It is surprising to find that a great number of physicians do not carry even the few items listed above, and are forced to use other less satisfactory preparations or call the "corner drug store" for supplies.

Chapter

I

ARRHYTHMIAS

A CHANGE in the rate or rhythm of the heart is a common cause of acute distress in patients with and without cardiac disease. Even in a patient with a normal heart an arrhythmia may produce many symptoms such as pounding, fright, precordial aching or severe pain,⁷¹ vertigo, syncope, vomiting, collapse, and rarely congestive heart failure.⁷⁰ The most severe circulatory derangements occur following the onset of ectopic arrhythmias with rapid rate, *i.e.*, the paroxysmal tachycardias.

PAROXYSMAL TACHYCARDIA

The paroxysmal tachycardias include auricular flutter, and fibrillation, auricular, nodal and ventricular tachycardia and ventricular fibrillation. These tachycardias often remit spontaneously within several hours and cause no disturbance, but they may produce severe symptoms and marked effects on circulatory dynamics. A change from a normal heart rate of 70 to 90 to one of 120 to 130 usually increases minute volume output for despite the fact that ventricular filling and stroke volume may be decreased, the product of heart rate and stroke output is increased. When the heart rate is between 120 to 130 and 180, however, no further increase in minute output occurs. With further increases in cardiac rate, total minute volume is actually reduced.⁷⁴ This may result in severe cerebral anoxia, fainting, or convulsions.

J C, a thirty two year old patient studied by us, was a tense person and a factory manager, who fainted on several occasions. He gave a long history of discomfort whenever he wore a tie and his pulse slowed moderately on carotid sinus pressure. The fainting was therefore considered to be the result of carotid sinus sensitivity. Close questioning however revealed that the syncope was sometimes preceded by palpitation. The patient was later seen during an episode of nodal tachycardia. This was doubtless the major cause of his fainting although the carotid sinus may also have played a role.

If the tachycardia is allowed to persist it may result in a further decrease in cardiac output, shock and/or cardiac failure.²⁰⁷ A picture of coronary insufficiency with chest pain and RS T and T wave changes in the electrocardiogram may also occur if the tachycardia continues. We have seen many such patients one of whom was a sixty five year old woman with carcinoma of the left lung and hemothorax who was receiving nitrogen mustard therapy. On the fifth day after therapy she suddenly experienced precordial pain and palpitation and went into shock. Her pulse rate was 180. A cardiogram revealed a 2:1 auricular flutter with RS T depressions in all leads. While the cardiogram was being taken the rhythm reverted to normal but the RS T depressions persisted for several minutes before disappearing. In many instances these RS T depressions persist for hours or days.

(The *supra ventricular tachycardias* i.e. auricular or nodal tachycardia and auricular flutter and fibrillation, may occur in both normal and abnormal hearts^{27, 143} and may be precipitated by emotional disturbances⁷². Occasionally, they are the only indication of hyperthyroidism; although a recent study has shown that the basal metabolic rate is within normal limits in over 85 per cent of all cases of supra ventricular tachycardia.¹⁴³ Most commonly, these arrhythm

mias occur in patients who have rheumatic or arteriosclerotic heart disease or in those whose hearts are normal Ventricular tachycardia on the other hand is almost always associated with organic heart disease particularly coronary occlusion¹⁵ and is of more serious import

/In many normal persons with paroxysmal tachycardia therapy may be withheld for several hours since the tachycardia often remits spontaneously. Individuals with normal hearts may be able to tolerate rapid heart rates for weeks or months without exhibiting any detectable alterations in cardiac function^{89 229}. Some patients however are emotionally disturbed and may experience precordial discomfort as a result of prolonged rapid beating of the heart in such patients treatment should be started early.

Certain general measures are of value in the prevention and treatment of all tachycardias namely rest sedation and the avoidance of tobacco alcohol caffeine and emotional upsets. A twenty eight year-old physician recently observed developed auricular fibrillation whenever he took one drink and smoked one cigarette simultaneously. He was finally persuaded to stop drinking and smoking and experienced no further attacks. Occasionally the ingestion of small amounts of alcohol will actually prevent the recurrence of tachycardias although this is not common. Eating fatty foods or large meals on rare occasions produces auricular arrhythmias in patients with otherwise normal hearts. We have observed a patient who suffered a bout of auricular tachycardia each time he ate a frankfurter or a large dish of ice cream. The avoidance of these foods completely eliminated the attacks.

{ A correct diagnosis is important before specific therapy for paroxysmal tachycardia can be instituted. It should be remembered that any of the tachycardias may be transitory and it is best to wait several hours before beginning drug therapy except in cases where the patient is in shock or cardiac failure.

AURICULAR FIBRILLATION

Auricular fibrillation is the most common serious arrhythmia and occurs most frequently in patients with rheumatic and arteriosclerotic heart disease³. It may be seen in cases of Graves' disease⁵⁰ following operative procedures⁸⁵ in pneumonia and in other febrile illnesses. Both the paroxysmal and chronic types are seen occasionally in the absence of organic heart disease^{4, 9, 98}.

Auricular fibrillation has been attributed by some observers to rapid irregular circus movements initiated in the auricular muscle and perpetuated by the occurrence of local blocks¹⁻⁶ but this view has been challenged by many authors^{1, 2, 7-11} who believe that both auricular fibrillation and flutter are the result of stimuli arising in a single exciting ectopic focus. According to the latter opinion all auricular arrhythmias have a similar mechanism the only difference between them being the number and frequency of impulses arising from the single focus²¹⁻⁷.

The paroxysmal form of auricular fibrillation usually sets in abruptly with a ventricular rate between 120 and 160 and a totally irregular rhythm. A pulse deficit may be present. When heart failure is absent it is best to observe these patients for several hours at rest and with sedation. If the fibrillation persists 3 grains (2 gm) of quinidine should be given orally every two hours. The first dose may be used to test sensitivity for reactions will usually occur within two hours if the patient is allergic to the drug¹⁹. True sensitivity is unusual however and treatment of a patient who is critically ill should not be delayed. If there is no change in rhythm after four hours the dose is increased to 6 grains (4 gm) every two hours. It must be remembered that quinidine has relatively little cumulative effect and that peak plasma levels are reached after 4 or 5 adequate doses have been given⁹. If the same dose is administered thereafter the plasma level rises only slightly and it is therefore necessary to increase the dose again to achieve

an effect. The total daily or weekly dose is of much less importance in determining plasma concentration than is the amount of the individual dose^{2,3} as only traces of quinidine are found in the plasma twenty four hours after the last dose has been given. Occasionally it is necessary to raise the dose to 9 grains (6 gm) every two hours. If regular rhythm is not restored with these doses larger ones will rarely have the desired effect and if the arrhythmia persists after 24 to 36 hours the drug should be discontinued.

We have used quinidine intramuscularly during the past year with excellent results. A suitable preparation is a 20 per cent solution of quinidine sulfate in propylene glycol 3 grains (2 gm) per cc. The preparation is stable and the injection is rarely painful⁴. The dosage is the same as when given orally and there is no greater toxicity when the drug is given by this route. Intramuscular medication is especially valuable when oral administration cannot be tolerated. Quinidine hydrochloride is also satisfactory for intramuscular use. Intravenous quinidine which is now available as quinidine lactate in 10 cc ampules containing 65 gm is a dangerous drug and its use must be carefully controlled. There is little indication for the intravenous administration of quinidine as most cases will respond to intramuscular or oral medication.

If the fibrillation persists after a satisfactory trial with quinidine the patient should be digitalized (*vide infra*). In all recalcitrant cases it is essential to exclude the presence of hyperthyroidism or active rheumatic heart disease. In

quinidine therapy after the hyperthyroid state has been corrected⁵.

If quinidine restores sinus rhythm the dose is reduced gradually and a maintenance dose is instituted. Three to six grains (2 to 4 gm) 3 times a day may suffice but 6 to 9 grains (4 to 6 gm) 4 or more times a day may be re-

quired for long periods. When quinidine is given q i d the first dose is taken preferably early in the morning on awakening the second at midday the third in the evening and the fourth dose just before retiring in order to maintain adequate blood levels. In unusual instances the fourth dose must be given during the night.

An example of this type of case is that of a fifty nine year old white male who had severe rheumatic heart disease and auricular fibrillation and was treated with 3 gm (5 gr) of quinidine sulfate orally every two hours as outlined above. Regular rhythm was restored and he was maintained successfully on 3 gm 4 times a day awaking at 4 A M daily for the morning medication. Whenever this dose was omitted fibrillation recurred.

Recently 3 grains (2 gm) enscals of quinidine have been introduced to avoid giving a dose during the night. The drug in this form is more slowly absorbed allowing for a greater interval between doses.

If only one attack has occurred or when episodes are infrequent quinidine need be given only for one week. After the acute attack however if paroxysms occur frequently prophylactic quinidine should be continued indefinitely.

Occasionally the attacks respond to quinidine with increasing difficulty as they become more frequent. In such cases moderate doses of digitalis given in addition to the quinidine are sometimes beneficial. When the attacks recur with increasing frequency and do not readily respond to quinidine it may be more advisable to slow the ventricular rate with digitalis than to attempt to restore sinus rhythm with quinidine. A good example of this therapy is presented by a woman of forty four who had had rheumatic heart disease with mitral stenosis for twenty years. She had developed a cerebral embolus three years before. Since then she had experienced frequent auricular premature beats and episodes of auricular fibrillation. At first these readily responded to quinidine but the attacks

became very frequent and lasted several days in spite of large doses of this drug. The patient was much distressed during the attacks and developed great anxiety regarding their recurrence. In the next attack digitalis alone was given until the ventricular rate fell to 70. Since then she has been maintained on 5 mg digoxin daily, has remained in chronic auricular fibrillation but has been entirely comfortable.

Although recent studies have shown that adequate cardiac output can be maintained by many patients with chronic auricular fibrillation if they are adequately digitalized¹⁴⁰ some of them do not do so well during fibrillation as when their rhythm is regular.⁹ Not only is their activity restricted but there is the ever present danger that thrombi will form in the auricles of the fibrillating heart and that embolization will occur. For these reasons we believe that all attempts to restore sinus rhythm should be made before resorting to digitalis alone although this therapy may occasionally be satisfactory as illustrated in the above case.

When employed properly quinidine is a safe and effective drug. The harmful effects following its use have in general been overemphasized. Several premonitory signs and symptoms of overdosage in patients under quinidine treatment should be kept in mind. The most common of these may be grouped under the term *cinchonism*. They include tinnitus, impaired hearing, headache, blurring of vision, giddiness, nausea, vomiting, abdominal cramps and diarrhea. If the drug is given too rapidly by the intravenous route respiratory depression and convulsions similar to those seen in animals may result.⁹⁶ Rarely a true *idiosyncrasy* may be present. In such cases acute respiratory distress and circulatory collapse appear even after small doses. Serious skin eruptions^{2,4} occasionally are a manifestation of quinidine sensitivity and rarely thrombocytopenic purpura develops (*vide infra*).^{100, 21}

A dramatic instance of quinidine sensitivity was seen in I. K., a teacher of sixty-eight, who had had a coronary

quired for long periods. When quinidine is given q i d the first dose is taken preferably early in the morning on awakening the second at midday the third in the evening and the fourth dose just before retiring in order to maintain adequate blood levels. In unusual instances the fourth dose must be given during the night.

An example of this type of case is that of a fifty nine year old white male who had severe rheumatic heart disease and auricular fibrillation and was treated with 3 gm (5 gr) of quinidine sulfate orally every two hours as outlined above. Regular rhythm was restored and he was maintained successfully on 3 gm 4 times a day awaking at 4 A M daily for the morning medication. Whenever this dose was omitted fibrillation recurred.

Recently 3 grains (2 gm) enoseals of quinidine have been introduced to avoid giving a dose during the night. The drug in this form is more slowly absorbed allowing for a greater interval between doses.

If only one attack has occurred or when episodes are infrequent quinidine need be given only for one week. After the acute attack however if paroxysms occur frequently prophylactic quinidine should be continued indefinitely.

Occasionally the attacks respond to quinidine with increasing difficulty as they become more frequent. In such cases moderate doses of digitalis given in addition to the quinidine are sometimes beneficial. When the attacks recur with increasing frequency and do not readily respond to quinidine it may be more advisable to slow the ventricular rate with digitalis than to attempt to restore sinus rhythm with quinidine. A good example of this therapy is presented by a woman of forty four who had had rheumatic heart disease with mitral stenosis for twenty years. She had developed a cerebral embolus three years before. Since then she had experienced frequent auricular premature beats and episodes of auricular fibrillation. At first these readily responded to quinidine but the attacks

symptoms listed above become manifest, the drug should be discontinued. If the patient is desperately ill, however, because of an arrhythmia, it may be necessary to continue quinidine administration despite mild symptoms of toxicity. Even in these cases, however, if severe symptoms of toxicity occur, the drug should be discontinued and other medication given.

Some authors have stated that quinidine should not be used in the presence of complete atrio ventricular or bundle branch block²¹ but we have not hesitated to employ this drug in supraventricular arrhythmias with bundle branch block, and have not observed any serious reactions as a consequence of this therapy.

When quinidine is unobtainable, oral potassium (5 to 10 gm daily given every four hours) may convert auricular fibrillation to sinus rhythm¹⁷⁹. This therapy was employed during World War II with excellent results.

In the treatment of acute paroxysmal fibrillation, occasional good results with Pronestyl¹⁸⁷ (*vide infra*) have been reported. These results as yet, have not been confirmed. This drug, however, is not successful in converting a chronic fibrillation to regular sinus rhythm, although slowing of the ventricular rate does occur.

If heart failure is present during the paroxysm of auricular fibrillation, digitalis should be administered immediately. It increases the degree of A-V block and slows the ventricular rate, but does not affect the fibrillation. The measures suggested below for the treatment of heart failure should also be instituted. Digitalization can be accomplished in many ways with various preparations. It is advisable to learn the potentialities of one preparation for oral use and of another for intravenous administration.

If the patient is acutely ill and immediate digitalization is indicated, strophanthin (K) or Ouabain (G strophanthin) should be given intravenously. These drugs have a very brief latent period, and are usually effective in from five to fifteen minutes, with a maximum effect in thirty to one

occlusion thirteen years ago and who recently experienced ventricular premature beats. On several occasions he developed acute laryngeal edema after receiving 3 grains (2 gm) of quinidine twice a day. A single daily dose finally sufficed to prevent the premature beats and no further episode of edema occurred while the patient was on this dose. Another graphic demonstration of quinidine idiosyncrasy was that of S. G., a physician of fifty-six with essential hypertension and coronary artery disease. On several occasions during the years he had received several grams of quinidine for ventricular premature beats. Six months ago he had taken 3 grains (2 gm) of quinidine 6 times during a two-day period when he developed profuse ecchymoses in the skin and mucous membranes and profuse hematuria. A blood count showed that the platelets had almost disappeared from the blood. The quinidine was discontinued and recovery occurred spontaneously within three weeks. These sensitivity reactions should be treated by immediate withdrawal of the drug. Blood transfusions should be given if required and antihistaminic preparations may be of great value.

Quinidine produces significant electrocardiographic changes, widening of the QT interval being the most common finding. This is due to widening of the R-T segment and only occasionally to widening of the QRS complex.¹⁰ The latter indicates a toxic effect of quinidine. These changes were noted in a thirty-four-year-old man who was admitted to the hospital because of recurrent episodes of palpitation and was discovered to have a nodal tachycardia. He was put on quinidine 6 grains (4 gm) every two hours. After 8 doses an electrocardiogram showed a QT interval of .58 seconds, largely as a result of widening of the R-T segment. When the quinidine was stopped the QT interval gradually returned to within normal limits within twenty-four hours. The QT interval may be even more prolonged than in the case cited above.¹¹ If significant widening of the QT interval is found or if some of the

symptoms listed above become manifest the drug should be discontinued. If the patient is desperately ill however

the drug should be discontinued and other medication given.

Some authors have stated that quinidine should not be used in the presence of complete atrio ventricular or bundle branch block⁴ but we have not hesitated to employ this drug in supraventricular arrhythmias with bundle branch block and have not observed any serious reactions as a consequence of this therapy.

When quinidine is unobtainable oral potassium (5 to 10 gm daily given every four hours) may convert auricular fibrillation to sinus rhythm.¹⁹ This therapy was employed during World War II with excellent results.

In the treatment of acute paroxysmal fibrillation occasional good results with Pronestyl²⁰ (*vide infra*) have been reported. These results as yet have not been confirmed. This drug however is not successful in converting a chronic fibrillation to regular sinus rhythm although slowing of the ventricular rate does occur.

If heart failure is present during the paroxysm of auricular fibrillation digitalis should be administered immediately. It increases the degree of A-V block and slows the ventricular rate but does not affect the fibrillation. The measures suggested below for the treatment of heart failure should also be instituted. Digitalization can be accomplished in many ways with various preparations. It is advisable to learn the potentialities of one preparation for oral use and of another for intravenous administration.

If the patient is acutely ill and immediate digitalization is indicated strophanthin (K) or Ouabain (G strophanthin) should be given intravenously. These drugs have a very brief latent period and are usually effective in from five to fifteen minutes with a maximum effect in thirty to one

hundred twenty minutes^{66, 14} A safe initial dose is 0.3 mg, within half an hour 0.1 mg may be given and repeated every half hour or hour until the ventricular rate is slowed to 10 or less¹⁵ Strophanthin K is available in 1 cc ampules containing 0.65 mg and Ouabain in 0.5 cc ampules containing 0.1 mg and 2 cc ampules containing 0.5 mg We have used both of these drugs intravenously with good results

Lanatoside C (Cedilanid) may also be given intravenously It has a slightly longer latent period and causes a decrease in heart rate in from twenty minutes to two hours⁶⁷ Its effect lasts longer than that of Strophanthin K or G making this drug ideal if moderately quick and slightly prolonged action is required The initial dose given is approximately one half the digitalizing dose i.e. 8 mg (4 cc) which is followed by 4 mg (2 cc) every four to six hours as required Cedilanid is available in ampules of 2 cc containing 4 mg and in ampules of 4 cc containing 8 mg

It is important to remember that these preparations are eliminated fairly rapidly and that the digitalization effect will be lost if longer acting digitalis preparations are not administered as soon as practicable i.e., within twenty four hours In this way digitalization is maintained Another very satisfactory method to follow if the patient is able to take medication orally is to give an initial intravenous dose and at the same time begin oral digitalization The intravenous medication is not repeated and oral digitalis is continued thereby giving rapid effect and prolonged action In general it is rarely necessary to use intravenous preparations except in patients with severe heart failure and rapid fibrillation

For oral use we have found Digoxin a glycoside of *Digitalis lanata* very satisfactory It has the advantage of being a pure substance of definite and constant composition and unlike digitalis leaf it is almost completely absorbed from the gastrointestinal tract²² It is eliminated rapidly and is not as apt to cause cumulative toxic effects such as those

caused by more slowly excreted glycosides, *e g* , Digitoxin. The effect of a single dose of Digoxin usually persists only for several days,¹⁴ whereas the effect of Digitoxin may continue for two to three weeks.¹⁵ We believe that at the present time, Digoxin is a very safe and effective preparation for oral digitalization.

The amount of any digitalis preparation required for digitalization varies greatly from person to person and the dosage must be individualized. The digitalizing dose of Digoxin varies between 1.5 and 5.0 mg.¹⁶ An initial dose of 1.0 to 1.5 mg is given, thereafter, 0.5 mg is given every six hours until the desired effect is obtained. The maintenance dose is 0.25 to 0.75 mg (1 to 3 tablets) daily, most patients being well controlled on 5 mg. Frequently giving 25 mg and 50 mg on alternate days proves very satisfactory.

Many physicians still use the whole digitalis leaf preparation for both digitalization and maintenance. This preparation is a good one, but the delayed onset of its action and the long period required for its dissipation are usually disadvantageous. When the patient will not take medication regularly, however, the use of the leaf is preferable, since the effect of Digoxin disappears in forty-eight hours and the patient may go into cardiac failure if he neglects to take it for two or three days. This danger is less when digitalis leaf is used. The whole leaf is much safer than Digitoxin and usually gives good results. Initial digitalization with the leaf can be accomplished with a total dose of 1 to 1.5 gm (15 to 22½ grains) given over a period of twenty-four to forty-eight hours, as desired. A useful guide for dosage is to give 1 gm (gr 1½) per 10 pounds of body weight.¹⁶ Adequate maintenance is usually achieved with 1 to 3 grains (0.6 to 2 gm) daily.

The patient should be observed carefully for signs of digitalis toxicity. Not infrequently, the most important and earliest of these are abdominal distention, cramps or pain, and visual disturbances, such as yellow or green vision,

and scotomata⁷¹. Occasionally cerebral symptoms such as confusion, dizziness and delirium appear. Nausea, vomiting and diarrhea occur in most cases of digitalis over dosage but may be late manifestations¹⁰. The most significant sign of overdigitalization is an arrhythmia resulting commonly from ventricular premature beats which occur in pairs or triplets and produce bigeminal or trigeminal rhythm. Bradycardia and various degrees of A-V block may also occur. It must be emphasized that premature beats may be found in patients who are not nauseated and do not vomit and if they occur the drug should be discontinued or the dosage decreased.

Depression of the RS-T segment and or inversion of the T wave in the electrocardiogram indicates that digitalis has had an effect on the heart muscle but does not imply toxicity. In patients with hyperthyroidism digitalis is usually ineffective even when tremendous doses are given. Conversely when a patient with auricular fibrillation or congestive failure fails to respond to adequate doses of digitalis one should suspect the presence of hyperthyroidism. On the other hand some people with hyperthyroidism have an increased susceptibility to digitalis and readily develop toxic symptoms²⁸. Careful studies with radioactive iodine may reveal hyperthyroidism as the cause of certain instances of such digitalis sensitivity.

The various arrhythmias resulting from digitalis over dosage usually disappear spontaneously if Digoxin was used they disappear within two to three days after the drug has been discontinued but they may continue for two to three weeks if Digitoxin had been given. Usually no treatment is necessary for the arrhythmia except with drawal of the drug⁷. In some instances of supraventricular tachycardias⁴¹ ventricular premature beats or ventricular tachycardias¹⁰⁹ induced by digitalis specific treatment may become necessary. In these cases potassium chloride or acetate given orally usually eliminates the arrhythmias within thirty minutes^{77,80} (Fig. 1). Two to ten grams of a



A



B



C



D

10 1-5 W Hypertension and moderate disease Digoxin toxicity 1 Lead III 40 Ventricular premature beats per 100 beats B Lead II 15 minutes after 10 (m Potassium orally 6 premature beats per 100 beats C Lead II 30 minutes 8 premature beats per 100 beats 1 Lead II 45 minutes No premature beats (Courtesy of Dr Charles I. N. Ivers)

20 per cent solution may be given in syrup of citric acid or smaller quantities may be given in milk, orange juice or ginger ale. When potassium is given in this manner nausea and vomiting usually do not occur. The effect produced will only last for from two to four hours and the treatment may have to be repeated. Toxic effects of potassium do not occur if there is adequate urinary excretion. Rarely is it necessary to use potassium intravenously; if it is, extreme caution should be used and only 5 to 15 gm. should be given slowly. Electrocardiograms should be taken frequently to detect potassium toxicity. Marked peaking of the T waves, widening of the QRS complex and disappearance of P waves are the most commonly observed changes when potassium intoxication occurs.¹⁷

Atropin is often effective in abolishing the A-V block caused by digitalis.

In auricular fibrillation secondary to hyperthyroidism or active rheumatic fever digitalis is ineffective. After the hyperthyroidism has been treated with radioactive iodine or by other methods the arrhythmia often disappears spontaneously¹⁸ or responds to quinidine therapy.¹⁹ A case recently studied illustrates this point. A fifty-three year-old female complained of constant palpitation. She had been found to have auricular fibrillation and had received very large doses of digitalis and quinidine without abolition of the arrhythmia or slowing of the ventricular rate. Although clinical signs of Graves' disease were absent, only 20 per cent of a test dose of radioactive iodine (I_{131}) was excreted and a diagnosis of hyperthyroidism was made. A therapeutic dose of 5 millicuries of I_{131} was administered. Two days later she was started on 30 drops of Lugol's solution daily. This medication was continued for four weeks and while the fibrillation persisted the ventricular rate fell to 84. The arrhythmia was still present six weeks after the administration of I_{131} and for this reason 6 grains of oral quinidine were given every three hours. Sinus rhythm ap

peared after 8 doses and persisted even after quinidine was discontinued

Radioactive iodine is usually picked up by the thyroid gland within twenty four hours but its effect on metabolic processes may not become evident for four to six weeks²⁶ It may be necessary therefore as in the case cited to administer Lugol's solution (10 drops t i d) during this period This medication may be started twenty four hours after radioactive iodine has been given and should be continued for approximately three weeks²⁷ The effect of the Lugol's solution will last for another week at which time the radioactive I_{131} effect will have begun to manifest itself

¹ If propyl thiouracil alone is used in hyperthyroid patients an effect on the arrhythmia should be seen within two to four weeks²⁸ It should be emphasized that the prognosis in these cases of auricular fibrillation due to hyperthyroidism is excellent if the primary disease is controlled¹⁷

Cases of auricular fibrillation occurring during active rheumatic fever usually respond to digitalis or quinidine therapy when the active process has been controlled Failure to respond to digitalis suggests that the underlying disease is still active The prognosis in these cases is not nearly as good as in patients with hyperthyroidism especially if a mitral valvular lesion is present²⁹

AURICULAR FLUTTER

¹ In auricular flutter, the ventricular rhythm is usually regular with a rate of 125 to 180 but it may be irregular because of varying degrees of A V block and thus may resemble auricular fibrillation Auricular flutter may sometimes be distinguished from fibrillation by the observation of rapid regular venous pulsations in the neck at a rate of over 250 corresponding to the rate of the flutter waves in the electrocardiogram Usually the atrial rate varies between 250 and 350 and a 2:1 A V block is present When auricular flutter is complicated by a bundle branch block

pattern it may simulate ventricular tachycardia electrocardiographically (Fig 2). Occasionally auricular flutter can be diagnosed with certainty only in the electrocardiographic leads taken to the right of the sternum³ or from the right chest or back. In rare instances it is necessary to take esophageal leads to record the auricular waves.⁴




FIG 2—S.W. m 71. Auricular flutter treated with intramuscular quinidine 0.6 Gm and oral Digoxin 0.8 mg. 6/7 S.P.M. 2/1 auricular flutter. 6/9 P.M. 1 hour after therapy regular sinus rhythm. 7/2 2/1 auricular flutter with bundle branch block simulating ventricular tachycardia.

Auricular flutter is usually paroxysmal but it may become chronic and continue for months or years.¹⁰ If a paroxysm does not remit spontaneously after several hours an attempt should be made to stop the attack with digitalis

or quinidine (Fig 2) digitalis being employed in all cases with heart failure. Digitalis slows the ventricular rate and increases the efficiency of the heart.

The fact that digitalis will oftentimes convert auricular flutter to fibrillation and that restoration of sinus rhythm will occur when the drug is discontinued has been repeatedly stressed¹⁶ since the original studies in 1911.¹⁵⁷ Although recent observations²⁹ have cast some doubt upon this theory we have seen a great many patients follow this pattern when digitalis is administered. //It has been our policy to administer digitalis convert flutter to fibrillation and then stop the digitalis in the hope that sinus rhythm will result. If sinus rhythm does not occur quinidine is employed to restore the normal rhythm. Usually a small dose of quinidine will be effective. Other observers also favor this approach to the therapy of auricular flutter.^{250, 266} Digitalis and quinidine are employed as described under auricular fibrillation. In critically ill patients there should be no hesitation about using intravenous digitalis followed by intramuscular quinidine if necessary.

In some instances auricular flutter persists in spite of all these measures and the patient must be maintained on digitalis as in chronic auricular fibrillation in order to keep the ventricular rate between 65 and 75 beats per minute. Some patients may continue to have auricular flutter for years and then suddenly remit spontaneously.²⁸⁵

If attacks of paroxysmal auricular flutter recur frequently an attempt should be made to prevent them by the continuous use of quinidine. The dose of quinidine for maintenance is 3 to 6 grains (0.2 to 0.4 gm) 4 times a day. In some cases however we have found it necessary to increase the dose and the frequency of administration giving as much as 9 grains (6 gm) 4 or 5 times a day. If this is not effective digitalization should be carried out and a maintenance dose continued. In this way a certain degree of A-V block is maintained and when an attack occurs the ventricular rate may not markedly increase despite a rapid

auricular rate. This may serve to keep the patient symptom free during an attack. This method is not successful in all cases for, despite adequate digitalization, a tachycardia may develop and symptoms appear.

AURICULAR AND NODAL TACHYCARDIA

Auricular and nodal tachycardia cannot be distinguished clinically and are treated alike. These tachycardias are frequently functional and may be precipitated by tobacco, alcohol, infections, gastric distention, Graves disease, allergic reactions and tense emotional states. Very frequently there is a combination of causes. The attack generally begins suddenly although in certain instances there may be a short premonitory period. The ventricular rate usually is regular, despite changes of position or exercise,⁷⁹ and is between 180 and 220 beats per minute, which is somewhat more rapid than in auricular flutter. Although there is usually a 1:1 ventricular response, occasionally a 3:1 or 2:1 response occurs with a resultant ventricular rate between 75 and 100.⁸⁰ These cases are more common than was previously suspected (Fig. 3). Frequently a supraventricular tachycardia can be differentiated from auricular flutter by the absence of flutter waves in the cervical veins. Unlike cases of auricular flutter, in which only transient slowing of the ventricular rate may be noted following vagal stimulation, supraventricular tachycardia often is terminated by this procedure. Supraventricular tachycardias are particularly apt to be evanescent, and patients may have many attacks daily for years without experiencing difficulty. Patients subject to recurrent attacks have learned, through experience, that certain maneuvers stimulating the vagus nerve abolish an attack, e.g., sudden movements of the head, holding the breath, coughing, vomiting, eating, lowering the head over the bed, or bending forward in a chair.

Often the attack disappears spontaneously but, if it persists, an attempt should be made to terminate it by




FIG. 3.—P E m 52 Coronary artery disease P roxysmal auricular tachycardia
190 ventricular rate 95

mediately when given by vein. Mecholyl should not be given to allergic or asthmatic patients because of the danger of inducing bronchial spasm or to hyperthyroid patients because auricular fibrillation may be precipitated.¹⁹⁴ It is best to keep the patient in a horizontal position when this drug is used to avoid possible syncope. Mecholyl and carotid sinus pressure usually are ineffective in cases of auricular tachycardia with A-V block and in auricular flutter and fibrillation and they may be ineffective if quinidine has been administered previously.⁴⁰

A less toxic parasympathomimetic drug which is effective in the treatment of supraventricular tachycardias is *acetylcholine*.²³¹ The effect of this drug is brief and its side effects are less marked than those of Mecholyl. Although we have had little experience with this preparation other observers have found it to be satisfactory when given in a single intravenous dose up to 100 mg. Certainly acetylcholine deserves a more extensive clinical trial.

Neostigmine, another parasympathomimetic drug is also effective in abolishing supraventricular tachycardias and is given in doses of 1 mg. intramuscularly.³⁴ When any one of these drugs is used atropine should be available for immediate use should it be necessary.

In cases in which mechanical methods are unsuccessful in which the distressing side effects of Mecholyl preclude its further use and in which more rapid action than is usually provided by quinidine is desired *neosynephrine* should be employed. When given intravenously in doses of 5 to 10 mg., this drug usually stops an attack of supraventricular tachycardia within twenty to thirty seconds.²⁷⁵ It produces a rise in blood pressure stimulation of the cardioinhibitor fibers in the aortic arch and carotid body and reflex cardiac slowing. Most attacks revert when the systolic blood pressure has risen to 160 mm. and the pressure usually returns to normal within ten minutes or less. There are few toxic effects of the drug but it should not be used in patients whose blood pressure is elevated during an attack or in




FIG 4—MB m 45 Paroxysmal nodal tachycardia of unknown etiology Cessation 10 min following 0.850 gm procaine amide intravenously Continuous lead II (Kayden Steele Mark and Brodie Courtesy of Circulation)

cases with ventricular premature beats. We have found neosynephrine effective in several cases after all other methods had failed.

As in the case of paroxysmal auricular fibrillation and flutter of recent origin Pronestyl occasionally stops auricular or nodal tachycardia.¹⁴⁴ Figure 4 shows the rapid effect of 0.8 gm. of the drug given intravenously to a man with nodal tachycardia. This patient of forty-five developed repeated episodes of this arrhythmia but no cardiac disease could be demonstrated.

It should be remembered that supraventricular tachycardias are so frequent and so often remit spontaneously that the accurate appraisal of the efficacy of any drug in their treatment is difficult. If there are numerous recurrences of the tachycardia the frequency of the attacks may be diminished by the continued use of quinidine or digitalis.

VENTRICULAR TACHYCARDIA

Ventricular tachycardia is infrequent and is usually associated with severe myocardial damage particularly that caused by coronary artery disease. It may be seen in cases of rheumatic heart disease during cardiac catheterization and in normal individuals.⁶ Occasionally it is possible to differentiate ventricular tachycardia clinically from auricular tachycardia by a slight irregularity in rhythm and a variation in the intensity of the first heart sound. Prominent jugular vein pulsations representing a summation of the auricular systolic wave (A wave) and the ventricular systolic wave (C wave) may be observed at a rate slower than that at the apex⁹¹ although this is not a common finding.

Digitalis is often a factor in the production of ventricular tachycardia.¹⁶⁵ Some observers believe that all cases of the bidirectional type of this arrhythmia are secondary to digitalis overdosage.¹⁶² We have stressed the fact that digitalis should be discontinued in all patients who develop ventricu-

lar premature beats especially if they are multifocal. If this precaution is taken many cases of ventricular tachycardia can be prevented.

Ventricular tachycardia usually responds readily to quinidine, given orally or intramuscularly occasionally an intravenous dose 1 gm (15 grains) has been required to stop an attack.²¹ We believe however that quinidine should be used intravenously only in dire emergencies because of its toxicity when given by this route and because better safer preparations are now available for the treatment of ventricular tachycardia.

A procaine derivative procaine amide, (*Pronestyl*) has been introduced for the treatment of ventricular arrhythmias.¹⁶⁷ This drug is a direct myocardial depressant. It is supplied in 250 mg capsules for oral use and in ampules containing 100 mg per cc for intravenous administration. We have used it with excellent results in ventricular tachycardia and premature beats including cases in which quinidine has failed. *Pronestyl* is given orally in doses of 500 mg every three to four hours or intravenously in doses of 250 mg to 1 gm at a rate of 100 mg per minute every thirty to sixty minutes until an effect is obtained. An electrocardiogram should be taken while *Pronestyl* is being given intravenously as the drug is often effective immediately. If a change in rhythm or widening of the QRS complex occurs the drug should be stopped. Occasionally oral doses up to 6 or 10 gm are required. We have found that both the intravenous and oral routes are satisfactory for the treatment of the acute episodes but the intravenous administration of *Pronestyl* produces a much more rapid effect. Ventricular tachycardia may be converted to regular sinus rhythm in from thirty seconds to two minutes following as small a dose as 250 mg of *Pronestyl*. A case recently observed is of interest to illustrate the effect of *Pronestyl*.

A sixty six year old white male experienced an attack of squeezing severe chest pain radiating to the left shoulder and arm, ten days prior to hospital admission. Four days

prior to admission he developed dyspnea on the day of admission he became dizzy and weak noted palpitations and fainted. On admission he was in acute pulmonary edema and was given 3 mg strophanthin K intravenously




FIG 5-NY m 66 Acute Coronary Occlusion Precordial pain heart failure and occasional palpitation on 10 days Strophanthin K 0.4 mg just prior to admission Irons 10 gm orally 3:45 P Dextro 10 mg orally at 6 P 1

morphine and 2 cc Mercurhydriⁿ intramuscularly. The electrocardiogram showed runs of ventricular tachycardia and acute anterior wall infarction (Fig 5). He was given 1 gram of Pronestyl by mouth and normal sinus rhythm developed within an hour (Fig 5). The Pronestyl was continued in doses of 1 gram every four hours for two days and then in doses of $\frac{1}{2}$ gram every four hours for eight days. He was also digitalized with Digoxin kept on maintenance doses and made an uneventful recovery.

Although Pronestyl generally does not produce serious reactions it may cause a marked fall in blood pressure or even cardiac standstill when it is administered intravenously. This may be prevented or counteracted by Neosynephrin[†] (1 to 5 mg) intramuscularly. The injection should be given slowly and blood pressure readings should be made during the injection. When the drug is given orally in large doses it may produce nausea and other gastric symptoms making it necessary to stop the drug or reduce the dose. We observed one patient who developed agranulocytosis while receiving Pronestyl and this has recently been pointed out by others. Therefore frequent blood counts should be made and evidence of purpura or bleeding watched for.

Pronestyl is also invaluable for the prevention and treatment of ventricular arrhythmias associated with operative procedures¹⁴ and may be of value in intracardiac catheterization. In preparation for the latter procedure it can be given prophylactically and if necessary during and after the manipulation. If an arrhythmia occurs during cardiac catheterization despite this therapy the catheter should be partially withdrawn and should not be advanced until the arrhythmia disappears. If used preoperatively 1 to 2 gms are given orally forty five to sixty minutes prior to surgery and 1 gm is given intravenously as the incision is made. During anesthesia and operation intravenous procaine alone may be used in a dose of 5 cc of a 1 to 2 per cent solution to prevent ventricular arrhythmias.⁴¹

Pronestyl has now been used prophylactically by us in a small series of patients who had previously experienced repeated bouts of ventricular tachycardia. Although this drug is able to terminate the acute attacks quickly and effectively, it does not appear to be completely effective in preventing recurrence of attacks when used orally 3 to 4 times daily. It may be that the dose employed (500 mg t i d or q i d) is not adequate and patients with repeated attacks should perhaps be given much larger doses. At the present time the best procedure appears to be to give Pronestyl for the acute episode and quinidine prophylactically.

Many other drugs that have been used in the treatment of ventricular tachycardias deserve mention at this time. One of these agents, intravenous *magnesium sulfate* is not widely used but has proven effective in occasional cases.²³⁻²⁶ When given rapidly in 2 to 4 gm doses i e 10 to 20 cc of a 20 per cent solution, this drug may terminate an attack after other measures have failed. Transient unpleasant side effects such as nausea, flushing, weakness, and dizziness may appear. It should be reserved for use after other therapy has proven ineffective. With the newer agents available, little opportunity for the use of $MgSO_4$ will present itself. Atabrine in doses of 4 gram intramuscularly may occasionally be effective.²³ Intravenous morphine sulfate 10 to 20 mg ($1/6$ to $1/3$ gr) repeated every hour if necessary, has been used successfully by several different observers.²⁹⁻³¹ The use of potassium salts (citrate or acetate in syrup or water, or chloride as enteric coated tablets) orally either alone in doses of 1 to 5 gms 2 to 4 times daily or in conjunction with quinidine has recently been advocated for the treatment of refractory cases or to diminish the frequency of recurrences of ventricular tachycardia.^{24,32} We have found potassium to be particularly useful in the treatment of ventricular tachycardia secondary to digitalis overdosage and in patients who are sensitive to quinidine.

Dibenamine hydrochloride an effective adrenergic blocking agent has not as yet been used by us in the treatment of tachycardias. This drug has been shown to be most effective in blocking arrhythmias which occur during cyclopropane anesthesia ^{121 128 129} and deserves a clinical trial in ventricular tachycardia which is not precipitated by operative procedures.

VENTRICULAR FIBRILLATION

Ventricular fibrillation is rarely diagnosed clinically and although it probably accounts for some cases of sudden death it is difficult to state its exact incidence. It occurs particularly in association with myocardial infarction, anesthesia, or excessive amounts of digitalis. The patient becomes pulseless and unconscious and rapidly goes into shock. Convulsions occur if this arrhythmia persists for more than thirty seconds. The electrocardiogram shows bizarre ventricular complexes and a completely irregular rhythm. Clinically, these attacks resemble a Stokes Adams seizure secondary to complete heart block and cardiac standstill and it is important to distinguish between these two conditions electrocardiographically. The use of sympathomimetic drugs *ie* epinephrine or ephedrine so valuable in the treatment of cardiac asystole is contraindicated in ventricular fibrillation. If possible the physician should obtain an electrocardiogram during the attack to establish its exact mechanism. Once an episode is diagnosed correctly as ventricular fibrillation it can be assumed that subsequent attacks will be caused by a similar mechanism and not by cardiac standstill and they may be treated accordingly. Bouts of both ventricular fibrillation and cardiac asystole may however occur in the same patient (Fig 6).

Since an immediate effect upon the arrhythmia is necessary, intravenous quinidine should be given promptly as quinidine lactate 4 to 6 gm (6 to 9 gr) despite the fact that the use of this route of administration is not without danger. Magnesium sulfate, 10 cc of a 20 per cent solution, may also be effec





FIG 6 - M V m 60 Coronary artery disease Complete A V block with periods of ventricular fibrillation and Stokes-Adams seizures Lead II continuous

tive. Other drugs have been suggested for the treatment of this condition but have had limited trial. Intravenous dibenamine has been shown to be effective in preventing ventricular fibrillation in animals and may prove to be of value in human beings. Intravenous atropine, 2 mg (gr 1/30) has been used successfully to abort the acute attack and smaller doses .5 mg (gr 1/100) intramuscularly have been given daily to prevent recurrent bouts in patients subject to paroxysmal ventricular fibrillation.²⁷ Intravenous Pronestyl has not been used by us for this condition but it should prove to be of value. In many cases the attack will cease or the patient expire before therapy can be instituted.

HEART BLOCK

Complete heart block with sudden failure of the idioventricular pacemaker and asystole is the commonest cause of Stokes Adams syndrome although a similar clinical picture may be seen in cases of ventricular tachycardia or other arrhythmias.²⁰³ Heart block is usually seen in patients with organic heart disease secondary to coronary arteriosclerosis but a small percentage of cases results from digitalis overdosage.¹² Congenital heart block is much more common than was previously believed and may account for 10 per cent of all cases.^{49, 5} Many patients experience only momentary weakness or unconsciousness and require no therapy but others have severe symptoms and require immediate treatment. One half to one cc of a 1/1000 solution of epinephrine is given subcutaneously or directly into the heart in extreme cases in shock. If the seizures recur the Adrenalin should be repeated subcutaneously every hour or two or may be given in oil (1 cc = 2 mg) intramuscularly every twelve to twenty four hours. We have observed a fifty six year-old female who required 5 cc of Adrenalin every fifteen to thirty minutes for hours in order to remain free of attacks. Intracardiac Adrenalin was necessary on one occasion after intramuscular therapy.

had failed and during a period when the patient experienced recurrent Stokes Adams seizures a slow continuous intravenous injection of 1 100 000 Adrenalin at a rate of 2 0 cc (30 drops) per minute was maintained for several hours in order to stop the syncopal episodes In patients who experience frequent episodes of Stokes Adams seizures ephedrine gr $\frac{1}{4}$ to $\frac{1}{2}$ (15 to 30 mg) or paredrine gr $\frac{1}{4}$ to 1 (20 to 60 mg) may be given orally 3 to 5 times daily in an effort to decrease the number of recurrences ²⁷ In all cases where digitalis may be a contributing factor this drug should of course be withdrawn

OTHER ARRHYTHMIAS

Sinus bradycardia sinus tachycardia and premature beats may occasionally cause severe difficulty These will be discussed briefly

Sinus tachycardia is usually associated with febrile or toxic conditions congestive heart failure or emotional or metabolic disturbances There is no specific therapy other than that directed at the primary condition

Sinus bradycardia, like sinus tachycardia may be an incidental finding especially in athletes and pregnant women and usually requires no treatment It is caused by an increase in vagal tone or by a diminution in sympathetic tone or both ²⁴ If the rate falls below 40 dizziness and syncope may result as in a person with carotid sinus syndrome Actual cardiac standstill may occur ²⁰⁷ We have recently observed a basketball coach of thirty two who usually had a pulse rate of 48 beats per minute He fainted on several occasions during tense moments in a game During the attacks of syncope his pulse rate fell to 30 and he usually recovered after thirty to forty seconds Usually these cases respond to ephedrine in 25 to 45 mg (3 8 or $\frac{3}{4}$ gr) doses or to atropin sulphate (6 to 1 2 mg) gr 1/100 to gr 1/50 by mouth It may be necessary to give

atropin (6 mg to 2 mg) gr 1/100 to 1/30 subcutaneously or intravenously

Premature beats may cause palpitation precordial discomfort and anxiety particularly if they occur frequently. These symptoms can usually be relieved by reassurance rest and sedation but occasionally they are sufficiently severe and persistent to require active treatment. A definite relationship has been found between anxiety states and episodes of premature beats and it has been shown that they as well as the distressing symptoms which occur during attacks may in some cases be eliminated by adequate psychiatric therapy.²⁷² If drugs are necessary to stop premature beats the physician should determine the site of origin of the ectopic beats before beginning therapy. Quinidine is usually successful in reducing the frequency of both ventricular and auricular premature beats. Pronestyl (procaine amide) is most effective in stopping or preventing ventricular premature beats.

A forty two year old salesman began to experience frequent choking sensations associated with a momentary cessation of the heart beat and on several occasions he became faint and almost collapsed. Examination showed the heart to be normal but the electrocardiogram revealed ventricular premature beats. He received 250 mg of oral Pronestyl 4 times a day with prompt disappearance of the arrhythmia and has remained symptom free since that time.

Although this patient was controlled by 250 mgm of Pronestyl q i d most commonly 500 mg is required orally every four to six hours to attain a similar effect. Pronestyl is preferred to quinidine which was formerly employed and is more effective than is quinidine. Premature beats assume great clinical importance when they precede the onset of paroxysmal tachycardia especially during operative procedures or following a coronary occlusion.

Chapter

2

ACUTE PULMONARY EDEMA AND CONGESTIVE HEART FAILURE

MUCH uncertainty exists today concerning the pathologic physiology of heart failure and acute pulmonary edema since many previously accepted concepts of cardiovascular and renal physiology and of sodium metabolism have been discarded, and since new concepts have been introduced during the last ten years

Pathogenesis of Cardiac Failure — In the past the symptoms and signs of congestive failure were attributed to "backward" failure *i.e.*, myocardial insufficiency, (cardiac failure) with increased venous pressure and venous congestion, increased blood volume secondary to anoxia of the bone marrow and resulting edema.²¹⁹ However, recent investigations have shown that this mechanism does not explain many of the findings in congestive failure. Rather do they support the theory of 'forward' failure, *i.e.*, cardiac insufficiency, diminished cardiac output with diminished renal blood flow, and decreased sodium excretion, with resulting increase in blood volume and edema.¹⁸⁰ Most of the symptoms and signs found in congestive failure can be accounted for by these sequential events although they do not explain some of the findings in acute heart failure, *i.e.*, acute pulmonary edema (*vide infra*). Usually, when the cardiac output falls, there is a liberation of excessive amounts of antidiuretic hormone from the posterior pituitary gland²⁴ and of desoxycorticosterone from the adrenal cortex,²¹⁵ these, also, play a part in the retention of sodium and water, and contribute to edema formation. Although the patho-

genesis of congestive heart failure has not yet been completely clarified it is probable that elements of both the forward and backward hypotheses play a part in its development

Excellent reviews dealing with the mechanisms of cardiac failure have been written^{202 219 241} We wish however to emphasize certain facts presented in these papers concerning the patho physiology of the various types of heart failure A knowledge of these basic facts is essential for the intelligent handling of both the acute and chronic forms of this condition

Heart failure occurs when the myocardium is no longer able to pump sufficient blood to all the tissues to satisfy their metabolic requirements At first the diseased myocardium is able to respond to stresses by increasing the length of its muscle fibers and contracting more forcefully (*Starling's law of the heart*²³⁹) Eventually however the fibers are distended beyond physiological limits and are no longer able to maintain an adequate output for tissue needs

In congestive failure the cardiac output may be low or normal as in hypertensive arteriosclerotic or rheumatic heart disease or it may be normal or high as in thyrotoxicosis²⁴² anemia²⁴² beri beri⁴ Paget's disease⁴³ arteriovenous aneurysm⁴⁵ or chronic pulmonary disease²¹⁹ In these latter conditions although the output is not decreased it still is unable to meet the tissue needs²⁴³ A primary object in therapy therefore is to increase the cardiac output In cases of failure with low cardiac output this can often be done by decreasing the blood volume by means of salt restriction or phlebotomy The resultant diminished venous return and lowered venous pressure enable the overdistended inefficient cardiac muscle fibers to shorten to within physiological limits to contract with greater force and to increase stroke volume and output In cases of failure with high output correction of the primary condition usually brings relief Digitalis increases cardiac output in both types of heart failure although it is more effective in cases with low

output.⁴¹ In some cases of high output failure *i.e.* cor pulmonale, digitalis may even be harmful if given too rapidly. It probably acts directly on the heart muscle increasing the force of its contraction⁴⁶⁻⁴⁷ although a primary effect on the venous system has been suggested.⁶⁰ This observation has not been confirmed.

Sodium retention is probably the single most important factor in the production of the symptoms of congestive failure. This retention of sodium and the consequent increase in total blood volume has been attributed to diminished cardiac output, diminished renal blood flow¹⁹⁰ and decreased renal excretion of sodium and water.¹⁴⁶ A patient with heart disease may remain well compensated while his sodium is restricted, but may quickly go into cardiac failure if given salt, because the resultant marked increase in blood volume throws an additional load upon the already overburdened heart.¹⁹³ Adequate salt restriction with the use of mercurial diuretics to aid sodium excretion, is therefore, of prime importance in the management of congestive failure.

A marked increase in body activity increases the work of the heart. It may cause a patient who was well compensated during rest or mild activity, to go into severe cardiac failure. It is important, therefore, to regulate a patient's activity carefully in order to avoid sudden episodes of cardiac failure.

If the above principles are borne in mind in the course of treatment of congestive failure many lives will be saved and careless mistakes avoided, for example, phlebotomy is of great value in acute left heart failure secondary to hypertensive heart disease,¹²³ *i.e.* cardiac failure with a low output, but it may cause great harm if performed in other forms of heart failure *e.g.*, heart failure associated with chronic pulmonary disease, Graves' disease, or chronic anemia. In these latter conditions a high output is necessary to maintain adequate tissue oxygenation and measures that lower the blood volume, such as phlebotomy, should

not be used since they further decrease the amount of oxygen available to the tissues

Decreased renal blood flow and sodium retention account almost entirely for the symptoms and signs of chronic congestive heart failure: *i.e.* dyspnea, orthopnea, pulmonary congestion, distention of the neck veins, hepatomegaly, ascites, and peripheral edema. Additional factors must be sought, however, to explain the sudden occurrence of acute left heart failure, *i.e.* cardiac asthma and acute pulmonary edema

Paroxysmal Nocturnal Dyspnea—Many patients with mild left ventricular failure experience recurrent episodes of paroxysmal dyspnea several hours after they go to bed. They awaken suddenly and are acutely dyspneic; a few rales and pronounced wheezes over both chests appear, without other evidence of failure. The blood pressure is high and the pulse is strong. This paroxysmal nocturnal dyspnea, commonly called *cardiac asthma*, is probably the result of a redistribution of blood and an increase in the blood volume with hemodilution when the patient assumes the recumbent position. During the day blood is pooled in the splanchnic and peripheral areas, but when the patient lies down blood is redistributed to the thorax⁶⁴ which results in lung congestion. As a result of this congestion, nerve reflexes from the lungs are initiated, and bronchospasm, dyspnea and the other symptoms and signs of cardiac asthma result. The exact mechanism of this condition has not been completely clarified.)

Complete relief is often obtained by having the patient sit up or stand, or walk about if possible. Mild cases may be controlled by aminophyllin suppositories (5 gm or 7½ grains). Aminophyllin is safe when thus administered, and is completely absorbed. Large doses can be given without producing significant gastric upsets. In the more severe cases or in cases that do not respond to this simple measure, morphine sulfate should be given intramuscularly, 15 mg (¼ gr), or intravenously, 8 to 10 mg (1/8 to 1/6 gr)

when an immediate effect is desired. The intramuscular route is usually satisfactory. Morphine very effectively suppresses the pulmonary reflexes that serve to perpetuate symptoms and should be used liberally.

An attempt to decrease venous return by the use of tourniquets or bloodless phlebotomy should be made if morphine is ineffective. Peripheral resistance which is high in many of these cases because of sympathetic nerve discharge may be decreased by giving aminophyllin (0.5 gm intravenously)¹⁹. Digitalization may not be necessary in the milder forms of cardiac asthma. Recurrences can be prevented by means of a low sodium diet and mercurial diuretics and by having the patient sleep on three or four pillows or in a chair in the severe cases. If attacks recur in spite of these measures or if other signs of left and or right heart failure occur the patient should of course be digitalized.

Acute Pulmonary Edema The onset of acute pulmonary edema secondary to left ventricular failure may be extremely sudden with signs of circulatory collapse severe dyspnea and cyanosis. It may occur following exercise a large meal paroxysmal tachycardia a coronary occlusion trauma to the central nervous system or after severe emotional upsets. A discharge of impulses from the sympathetic nervous system has been suggested as a possible explanation in the latter conditions.¹⁶¹

A good example of pulmonary edema induced by neurogenic factors was recently provided by a 65 year old woman who had experienced occasional attacks of pulmonary edema three years before but had been quite well since then. One week ago she witnessed her husband fall and severely injure his scalp as a result of a cerebral hemorrhage. He quickly developed a left hemiplegia. His wife appeared to be taking all this well but about an hour later she suddenly began to wheeze and numerous rales appeared in both lungs. She was given Demerol 50 mg subcutaneously and gradually improved.

Occasionally pulmonary edema may complicate bulbar

poliomyelitis and for this reason intravenous hydration should be carefully supervised in these cases^{118, 961} The heart sounds may be obscured by numerous bubbling rales in the lungs usually numerous wheezes are also present In many cases the right heart is strained by the failing left ventricle and the pulmonic second sound is accentuated If the patient is seen early the blood pressure and pulse volume may be maintained A short time after the onset of symptoms however there is a marked fall in blood pressure and the pulse becomes rapid and thready i e shock occurs due to acute left ventricular failure *Morphine sulfate* should be given intramuscularly at once or intravenously in cases with peripheral collapse in which little or no absorption would otherwise occur The doses employed are similar to those given in cardiac asthma This drug usually produces a dramatic improvement but if no effect is noted it should be repeated in fifteen to twenty minutes Dilaudid 1.5 to 3 mg (1/40 to 1/20 grain) or Pantopon 20 to 30 mg (1/3 to 1/2 grain) may be used instead of morphine especially if the latter induces vomiting Although some physicians use atropine 0.4 to 0.6 mg (1/50 to 1/100 gr) routinely in the treatment of acute pulmonary edema we do not feel that it is of value in cases in which tachycardia is present it may actually be harmful

Despite the fact that the arterial blood is fully oxygenated in many cases Oxygen is of value in pulmonary edema⁹ Often when given by the meter mask under pressure (3 to 6 cm) it produces dramatic results A thirty year old negress para iii with rheumatic heart disease experienced mild toxemia of pregnancy She was delivered of a full term normal infant after a short labor but bled profusely and rapidly went into shock She received 500 cc of plasma and 2500 cc of blood within the next three hours after having lost an estimated 2000 cc of blood She recovered from shock but developed a marked pulmonary edema one hour later Despite repeated doses of morphine oxygen by tent 8 mg (4 cc) Lanatocid C and continuous bloodless phlebotomy

with tourniquets (*vide infra*) she did not improve. After one hour of vigorous therapy oxygen under pressure (4 cm) was administered by the anesthetist and within 10 to fifteen minutes marked improvement occurred. The rales gradually disappeared and the patient recovered.

To prevent recurrences oxygen may be given in this way for a period of fifteen minutes every few hours for the following twelve hours. Oxygen by nasal catheter usually is not satisfactory but an oxygen tent often is of value in warm weather. It should be remembered that the percentage of oxygen inspired by a patient when a nasal catheter or tent is used is never as great as that inspired through a mask (35 to 65 per cent as compared to 50 to 100 per cent). There is some evidence that 100 per cent oxygen may be irritating to the lung parenchyma particularly if taken for hours or days for this reason the use of a lower concentration is advisable.⁶

Intravenous aminophyllin 0.5 gm ($7\frac{1}{2}$ gr) is of particular value in the treatment of acute pulmonary edema when Cheyne Stokes respiration is present. When given by vein it should be administered very slowly the total dosage being injected in forty five to sixty seconds. In less serious cases the same amount of aminophyllin may be given by suppository.

If the pulmonary edema persists after the above measures have been tried 25 mg of Strophanthin K or Ouabain (Strophanthin G) and 2 cc of a mercurial diuretic Mercurhydrin or Mercupurin should be given intravenously in the same syringe. Thereafter the Strophanthin is repeated or Digoxin is used as outlined previously. Mercurial diuretics are more toxic when used intravenously³⁶ and should be given by this route only in acute emergencies the intramuscular or subcutaneous route should be employed as soon as possible thereafter. *Bloodless phlebotomy* by tourniquets³⁶ if employed properly often is markedly effective in patients with pulmonary edema secondary to low output cardiac failure.³¹ Tourniquets having been

placed on three extremities sufficient pressure is exerted to occlude the veins but not the arteries. Pressure is applied for fifteen to twenty minutes and then the tourniquets are released and rotated so that one extremity is always free of pressure. If after forty five or sixty minutes no definite improvement occurs a phlebotomy should be attempted 500 to 800 cc of blood being withdrawn rapidly. Withdrawal of less than 500 cc rarely produces the desired decrease in blood volume but if an adequate amount is withdrawn quickly a resultant fall in venous pressure and a rise in cardiac output can be expected. This increase in output occurs because the heart is once again able to respond to stress i.e. the decreased venous return allows the overdistended ineffective cardiac muscle fibers to decrease in size and contract more forcefully within the limits of Starling's law of the heart. The hemoglobin and hematocrit should be checked before phlebotomy is done and if the results obtained are markedly reduced from the normal values expected this procedure should not be considered. On the contrary in pulmonary edema secondary to severe anemia with high output failure small transfusions of packed red blood cells may be life saving. Actual phlebotomy should be done only in unequivocal cases of severe low output cardiac failure after other measures including bloodless phlebotomy with tourniquets have failed.

Therapy of Congestive Heart Failure — After the acute attack of left ventricular failure is over the usual management of congestive heart failure is instituted. Maintenance digitalization with Digoxin 1 to 3 tablets (25 to 75 mg) per day a low sodium diet and the judicious use of mercurial diuretics are the three cardinal features of long term therapy. In addition *penicillin* (300 000 to 600 000 u daily) in a single dose should be administered in cases of moderate or severe heart failure to prevent secondary lung infections and Dicumarol should be given to prevent pulmonary infarction especially in patients who will be

bedridden for long periods of time.²¹⁶ We believe that the immense value of these measures in congestive failure has been largely overlooked. In our experience they have improved the course of apparently hopeless cases.

Adequate restriction of salt can now be achieved while the patient enjoys a reasonably well balanced palatable diet.²¹⁷ During the first week after the acute episode of heart failure a diet containing not more than 1 gram of sodium should be prescribed. This can be accomplished by allowing only fresh fruits, unsalted vegetables, potatoes or rice, salt free milk (prepared from lanolac powder), cottage cheese, and salt free bread. If the patient does well on this regimen, small portions of meat and fresh water fish may be given. The patient should be warned against eating the following foods:

Salt butter, margarine or peanut butter

Crackers, cakes or pastries or any food made with baking soda

Smoked or salt cured meats and fish, ham, bacon, pork or corned beef

Canned soups or vegetables

Pretzels, salted nuts, potato chips, candy, beets, elvers, lima beans, spinach and sauerkraut

Olive pickles, catsup, mustard or salad dressings

A salt substitute, *i.e.*, potassium chloride, may be used but most patients find this preparation bitter and unsatisfactory and prefer to use no salt at all. Palatability may be increased by flavoring with pepper, vinegar, garlic, onions, vanilla, chocolate or cinnamon. There should be no restriction placed upon the ingestion of water. A fluid intake of between 2 and 3 liters per day seems to be the optimal intake for maximal sodium excretion.^{173, 179} Tea, coffee, fruit juices, ginger ale, lemonade, salt free milk (lanolac) and occasionally small quantities of alcoholic beverages may be taken. Frequent small meals are to be preferred to a few large ones and under no circumstances should a large meal be eaten within three to four hours before bedtime. The

patient should be cautioned about the use of sodium bicarbonate or laxatives containing sodium

If a salt poor diet *i.e.* a diet containing between 1 and 2 grams of sodium daily, is followed carefully, many recurrences of attacks of acute pulmonary edema or severe congestive failure can be eliminated

Patients who are obese should be put on a fairly rigid reducing diet, and their weight should be kept at a minimum, if possible. We have seen patients with severe cardiac disease and intractable heart failure who were greatly improved and well controlled after marked weight reduction. Obese cardiacs are much more difficult to control than patients with normal weight. Great stress, therefore, should be placed upon weight reduction as an integral part of the therapy of congestive failure. If these dietary precautions are observed patients may be able to continue for many months without any additional therapy. If however the patient is unable to adhere to a proper salt poor diet or if his weight increases or edema develops while on such a diet mercurial diuretics should be used as often as is deemed necessary to keep the patient edema free and at a stable weight.

Intramuscular Mercurhydrin in 1 to 2 cc doses has been used extensively, and is a safe effective mercurial which causes little pain at the site of injection. It may be used subcutaneously or orally but its action is not wholly satisfactory when given by these routes.

Thiomerin a mercurial which is given subcutaneously is effective and easy to administer,¹⁰⁵ and we have used this drug with excellent results. It is possible to teach patients to administer it to themselves, this greatly facilitates therapy and is as effective and safe as is insulin administration. Since exercise increases sodium retention,¹⁰⁶ patients are instructed to stay in bed, if possible, for an hour or two following each injection. In this way the maximum diuretic effect may be obtained. Occasionally minor muscle

cramps occur two hours after the first dose of this medication but this discomfort disappears quickly

Oral mercurial diuretics may produce a satisfactory response in patients with congestive heart failure²⁷ but the use of these preparations has thus far been limited because of toxic reactions²⁸ and uncertain effects. In patients who require two or more injections of a diuretic weekly it is often possible to eliminate one of the injections by using a longer acting oral mercurial (Mercurhydrin Mercurhydrin ascorbic acid Salyrgan or Mercuzanthin tablets) but usually the oral medication is not dependable enough to allow elimination of parenteral therapy. Occasionally however a patient can be completely controlled by oral therapy. A seventy one year old woman with chronic rheumatic heart disease with auricular fibrillation and with chronic heart failure was treated with Digoxin and mercurial injections for many years but because the mercurial injections proved painful they were discontinued. An oral mercurial preparation (Salyrgan theophylline) was then given (2 tablets daily equivalent to approximately 1 cc of the parenteral preparation). The dosage was reduced to 1 tablet daily after one year and the patient has been controlled satisfactorily on this medication for three years. Certainly if an oral mercurial preparation that produces a satisfactory prolonged effect is found it will further simplify diuretic therapy.

Occasionally it is necessary to use ammonium chloride 4 to 6 grams (60 to 90 gr) a day for two to three days prior to mercurial administration to potentiate the mercurial effect.²⁹ Enteric coated tablets containing 1 gm each given in small and frequent doses are well tolerated but the drug should not be used continuously or in patients with marked renal impairment because of the possibility of inducing severe electrolyte disturbances i.e. hypopotassemia and acidosis. Aminophyllin 0.5 gm ($\frac{1}{2}$ gr) given intravenously increases renal blood flow and when used sixty

to ninety minutes after a mercurial diuretic may markedly increase the diuretic response¹⁶⁸

Mercurials may be given safely in the presence of renal disease unless oliguria or azotemia is present but great care must be taken when administering mercurial diuretics to elderly persons to avoid excessive diuresis which may result in salt depletion dehydration and azotemia¹² For this reason it is wise to check the blood urea and the sodium or chloride levels from time to time When salt depletion occurs it is often attended by serious chemical derangements and symptoms¹²⁷⁻³³ the most important of which are extreme weakness apathy disorientation stupor coma and rarely death and while the daily injection of a mercurial often is dramatically effective in young persons it may be dangerous for the elderly patient For example

A seventy year old man who had been in congestive failure for four years had received a mercurial injection of 2 cc every week during that period and apparently had been well controlled until three weeks prior to admission He was admitted to the hospital for treatment of severe right and left heart failure The specific gravity of the urine was 1.020 the blood urea nitrogen 15 mg per cent the urea clearance 54 per cent in two hours and the phenol sulfonphthalein excretion 35 per cent After receiving 2 cc of Mercupurin by vein daily for ten days he had lost 17 pounds and all signs of congestive failure had disappeared The urea nitrogen had risen to 61 mgm per cent however and the patient had become disoriented lethargic and hyperpneic He began to vomit and developed a tremor of the hands Urinary output dropped to 85 cc in twenty four hours and the blood urea nitrogen rose to 109 with a blood chloride level of 679 mg per cent An intravenous drip of glucose in saline was given continuously but he died sixteen days after the last mercurial injection

For older people only 1 cc of Mercurhydrin should be given intramuscularly or 1 cc of Thiomerin subcutaneously not more than every other day after the acute attack of

heart failure has been controlled. In elderly men mercurials often aggravate prostatic disease causing acute retention and severe uremia^{279, 284}. In these cases mercurials should be given with even greater caution. It has been demonstrated that morphine, Demerol, quinine, ephedrine and Adrenalin often decrease the effectiveness of the mercurial diuretics and these drugs should be avoided if possible while the mercurials are being given^{81, 181}. It may also be wise to reduce the dose of digitalis on the day that a mercurial diuretic is given in certain patients who are markedly edematous. If this is not done, mobilization of digitalis with the edema fluid²⁸⁵ may result in the release of excessive amounts of digitalis and digitalis intoxication^{6, 7}. This complication of mercurial administration is not stressed but is probably much more common than previously suspected. We have seen several patients whose digitalis dosage had been well regulated and who developed signs of digitalis toxicity: *i.e.* nausea, vomiting and ventricular premature beats within twelve to twenty-four hours after the initial dose of a mercurial diuretic.

The judicious use of the mercurial diuretics with adequate salt restriction will usually control cardiac edema but in some refractory patients who are not able to stay on a salt poor diet and who are perpetually in chronic congestive failure the use of *cation exchange resins* may be justified if given carefully under close supervision. These resins which are substances capable of binding ingested sodium within the intestinal tract and excreting it in the stools⁶ are now available for general use. Hitherto their administration was restricted because a great many pills had to be ingested to produce effects and excessive amounts of potassium were eliminated along with other electrolytes resulting in severe electrolyte disturbances. The new ammonium cation exchange resins are prepared in powder form have added potassium and may not present this difficulty^{1, 4}. The safety of even these preparations has not been proven conclusively and they should be used cautiously and only

when close supervision of the patient is possible. Blood electrolyte determinations should be done frequently, at least for the first few weeks following the onset of therapy. If hypopotassemia or rapid sodium depletion occurs the resin should be temporarily discontinued. Our experience thus far has indicated that if administered carefully these substances are of value in the therapy of congestive failure especially in the cases where patients do not remain on an adequately controlled salt poor diet.

In addition to the above measures the patient's activity should be carefully regulated. Short periods of bed rest should be taken if symptoms become severe or if heart failure appears to be progressing.

In heart failure secondary to hyperthyroidism pernicious anemia beriberi or surgical lesions the correction of the primary disease usually prevents its recurrence. In cardiac failure secondary to tachycardia treatment of the latter condition should be instituted as outlined above.

In patients with heart failure fluid may accumulate rapidly in the chest and cause acute distress. In these cases immediate and repeated thoracenteses give remarkable relief and permit recovery. Accumulation of fluid in the abdominal cavity also occurs but rarely does enough fluid collect to cause symptoms and require paracentesis. This has been especially true in recent years since the use of good salt poor diets and adequate mercurial therapy became widespread. Occasionally in patients with severe rheumatic heart disease and long standing congestive failure massive ascites will develop as a result of advanced cardiac cirrhosis. We have seen patients who required frequent paracenteses in order to remain comfortable. One of these was a thirty four year old female who was hospitalized because of chronic congestive heart failure, secondary to rheumatic heart disease. Bilateral pleural effusions and ascites were found. Despite adequate digitalization, marked salt restriction and the frequent use of mercurial diuretics, little improvement was seen after one week of therapy. A paracen-

tesis was performed and 7 liters of fluid were removed. A marked diuresis with dramatic clinical improvement occurred during the next four days. Another paracentesis was done one week later thereafter the patient was adequately controlled and required no more tapping.

Cardiac cirrhosis and liver disease in patients with chronic congestive failure is apparently much more common than previously suspected^{25, 14} and for this reason attention should be paid to the patient's intake of protein and vitamins. Because these patients are often on a strict diet there is a tendency to overlook simple measures such as the use of protein and vitamin supplements that might postpone the occurrence of severe liver disease and ascites.

During attacks of acute heart failure especially following coronary occlusion annoying and often serious symptoms may occur e.g. hiccup severe nausea vomiting and distention the latter symptoms occasionally result in death. It is often of great importance to stop these symptoms quickly or prevent their occurrence.

The excessive use of opiates should be avoided if possible and laxatives should be given freely to avoid constipation. In the presence of continuous narcotic medication two and even three times the usual dose of the laxative may be necessary. If therapy with laxatives such as Milk of Magnesia (30 to 60 cc) or Cascara sagrada (10 to 30 gr) is not successful occasional low gentle enemas may be used in conjunction with the above. Cathartics that act upon the large bowel such as Cascara sagrada or rhubarb are not to be given however in patients receiving mercurials because of the possibility of inducing severe colitis.

We have found intravenous pyridoxine (25 to 50 mg) of great value in treating the nausea and vomiting which complicates congestive failure. Other measures that we have found of value in treating these symptoms include sedation restriction of foods and fluids sips of ginger ale Dramamine (100 mg every four hours if retained by mouth) and the elimination of the possible causative agents such as digi-

tals morphine aminophyllin and ammonium chloride. Certainly there is no indication for the forcing of cold milk and fruit juices as is done so often since these tend to increase nausea and the nurse or patient's family should be instructed specifically against this practice. However milk which is boiled and then cooled to room temperature is usually well tolerated. The fruit itself instead of the juice is also satisfactory.

Intractable hiccup has been treated successfully with sedation, amyl nitrite, intravenous niacin, inhalation of 5 to 7 per cent CO_2 and 90 per cent oxygen for a few minutes at a time, ethyl chloride spray to the region over the diaphragm, inhalation of ether or intramuscular quinine gr. 6 (4 gm.) 3 times a day.²³ A phrenicectomy or exeresis is occasionally required to stop a severe attack.

Chapter

3

ANGINA PECTORIS CORONARY INSUFFICIENCY AND CORONARY OCCLUSION TREATMENT

THE most common cardiac emergency is pain in the chest. It is essential to exclude extracardiac causes of chest pain such as spondylitis herpes zoster pneumothorax pleurisy gallbladder disease peptic ulcer hiatus hernia and esophageal spasm *before deciding that the pain is cardiac in origin*. This can usually be done by taking a careful history doing a complete physical examination and employing routine laboratory tests. Once the cardiac origin of the pain has been established the type of the acute cardiac episode must be determined. The character intensity and duration of the pain may help the physician to distinguish between attacks of coronary insufficiency including angina pectoris and coronary occlusion. The pain in coronary occlusion is prolonged severe and unrelieved by nitrites whereas in coronary insufficiency particularly angina pectoris it is usually briefer and is relieved by nitroglycerin. Occasionally however coronary occlusion is associated with only mild pain and no other symptoms whereas coronary insufficiency may cause severe pain. The entire clinical picture and the electrocardiographic changes should be considered before a definite diagnosis is made and occasionally it can be made only after observation for several days. Any patient therefore who experiences sudden precordial pain diagnosed to be cardiac in origin should be put to bed and

treated as a case of potential infarction although many of these cases do not progress to a coronary occlusion with infarction

Angina pectoris, precipitated by effort exposure to cold, excitement or following large meals is a good example of temporary coronary insufficiency with ischemia of the sub endocardial layer but no myocardial necrosis The ordinary episode of angina pectoris is quickly relieved by rest or nitroglycerin a hypodermic tablet gr 1/200 to 1 150 being given under the tongue The patient should be instructed to take the nitroglycerin immediately if the pain is severe or if a specific activity such as eating or walking in cold weather is known to induce anginal pain the drug should be taken prior to this activity For example many patients experience pain only in the morning on the way to the subway or train a nitroglycerin tablet under the tongue just before leaving the house often prevents the attack This drug is safe and extremely effective and should be taken as often as necessary If nitroglycerin is not used and the pain persists the patient becomes frightened and anxious this emotional state serves to perpetuate the attacks The establishment of a vicious circle of pain anxiety and more pain makes therapy extremely difficult but liberal use of nitroglycerin and sedation often interrupts this seemingly endless chain There is apparently no danger in giving nitroglycerin as frequently as every hour Occasionally however it produces flushes intense throbbing and severe headaches or even syncope and in these instances it should be used with caution For example A forty eight year old waiter suddenly experienced substernal pain and was given gr 1 150 of nitroglycerin Severe throbbing in the head which lasted for several hours profuse perspiration and faintness occurred but the pain was not relieved During another attack of precordial pain several days later the patient was prevailed upon to test the effect of nitroglycerin in a minute dose of gr 1/400 but, again the same symptoms were produced Of course,

if several doses of nitroglycerin do not control the attack or if the pain lasts for more than twenty to thirty minutes the medication should be stopped and the patient put to bed and observed carefully for other signs of coronary occlusion. Treatment should then be instituted as described below.

Whiskey is occasionally effective in the treatment of angina and sometimes a patient prefers to take small sips of whiskey instead of nitroglycerin for the attacks. A few cases do not seem to respond to either nitroglycerin whiskey or to adequate sedation. These patients may continue to have frequent severe attacks of angina which at first appear only after exercise but later recur frequently at rest *i.e.* *status anginosus*. In many instances this condition heralds the onset of a coronary occlusion. When *status anginosus* occurs therefore the patient should be put to bed under adequate sedation and carefully observed for several days. The period of bed rest and sedation may alleviate the pain and allow the patient to resume at least mild activity upon arising. In those patients who do not show evidence of occlusion but who continue to have severe bouts of anginal pain other methods of treatment have been used and are being evaluated.

Many observers have used antithyroid drugs^{20, 21} and even thyroidectomy²⁰ for the treatment of severe angina hoping thereby to reduce cardiac work by lowering body metabolism. Good results have been reported in some cases after all other treatment has failed but these procedures do not appear warranted because of drug toxicity or surgical morbidity and mortality. More recently radio active iodine I_{131} has been employed to produce a hypothyroid state in angina patients with some success in intractable cases.²¹ This type of treatment is usually not required in cases of angina pectoris and we have rarely resorted to the use of antithyroid therapy in euthyroid patients. Occasionally when hyperthyroidism is present or aggravates a pre-existing anginal syndrome, correction of

the hypermetabolic state by radioactive iodine may completely relieve the anginal attacks. In older persons there may be no obvious signs of hyperthyroidism and only after a complete study can the condition be diagnosed.

A F. was a sixty seven year old Italian housewife who entered the hospital because of severe recurrent episodes of substernal pain which began suddenly two weeks before. Her blood pressure was 160/80 her rhythm was regular and a grade ++ apical systolic murmur was heard. The LCG showed deeply inverted I waves in the standard and precordial leads a diagnosis of coronary insufficiency was made. In spite of complete bed rest and large doses of sedatives she continued to experience frequent attacks of pain which were only partially relieved by nitroglycerin. During the course of a thorough study her basal metabolic rate was found to be +40 per cent and she excreted only 10 per cent of a test dose of I_{131} indicating hyperthyroidism. She was given a therapeutic dose of I_2 and Lugol's solution and after several weeks her anginal seizures began to subside and finally disappeared completely.

Papaverine has been advocated for the treatment of angina but careful studies have shown that in doses up to 800 mg by mouth daily it has no significant effect upon the anginal syndrome²⁴. The original claims that Alpha Tocopherol (Vitamin E) was of value in the treatment of angina pectoris have also been refuted²⁵. We have used both of these medications with poor results.

Khellin an extract from the seeds of an Egyptian plant is the most recent drug to be used in the treatment of angina. It dilates the coronary arteries when given intramuscularly in doses of 100 mg or orally in 40 to 60 mg doses 3 times daily. Some favorable reports on the efficacy of *Khellin*⁴ have not been confirmed²⁶. Our own experience has shown it to be of little value thus far in most of the patients who have received it gastro intestinal symptoms such as nausea and vomiting have developed.

For many years various surgical procedures have been

employed to relieve pain in cases of angina that responded poorly to nitroglycerin that recurred frequently and that required liberal use of narcotics for their control. Paravertebral injection of the upper five thoracic sympathetic ganglia and their rami communicantes with alcohol¹² has occasionally provided relief for patients¹⁷ who were not amenable to other types of treatment. This method or the injection of the stellate ganglion alone with ammonium sulfate²¹ appears to be achieving some satisfactory results in experienced hands. More recently good results have been reported from the use of various ganglionic blocking agents such as tetra-ethyl ammonium chloride especially in patients with status anginosus.¹⁸ There does not appear to be any justification at the present time for the use of surgical procedures that attempt to increase collateral coronary blood flow such as omental grafts,²⁴ implantation of irritating substances into the pericardial cavity²⁵ or the anastomosis of the aorta to the coronary sinus as has been advocated.¹⁹

We have been most successful in improving the condition of our anginal patients by the frequent unhesitating use of nitroglycerin by reassurance and by the judicious use of sedatives and narcotics. We believe that most patients should be treated by these means and that the use of other drugs or surgical procedures is only rarely justified. In only one instance has it been necessary for us to resort to surgical therapy in a case of angina pectoris during the past four years. In this case an upper thoracic sympathectomy was performed but the result was unsatisfactory. The patient continued to have precordial pain and in addition developed severe postoperative neuralgia.

Coronary Insufficiency—In recent years many acute coronary episodes have been observed which simulate coronary occlusion at their onset but do not resemble it electrocardiographically and in their subsequent clinical course.¹⁷⁶⁻¹⁷⁷ In these cases the ECG may show only RS-T depressions and/or T wave inversion which usually appear

soon after the attack although, in some cases, they may not become evident for several days or a week (Fig 7) Fever, leucocytosis, or elevated sedimentation rate may also be present, although, in the milder cases where myocardial ischemia without necrosis occurs, these findings are usually absent




FIG 7 —N G m 53 Acute coronary insufficiency following physical exertion (Master and Jaffe courtesy of Postgraduate Medicine)

When attacks of precordial pain occur spontaneously, at rest, it is often extremely difficult to distinguish between those that are caused by simple coronary insufficiency and those that will progress to coronary thrombosis. For example:

Mr. I F, a storekeeper, forty-four-years old, had been observed over a number of years because of numerous functional complaints. Physical examination, blood pressure,

and ECG were normal. One day while at work he experienced severe substernal pain which lasted one hour and closely resembled the pain of a coronary occlusion. At this time he was not in shock, his heart sounds were regular at a rate of 76 and of good quality. He showed diphasic T waves in leads I, II and V_{4-6} but his temperature was not elevated and there was no leucocytosis. He remained asymptomatic but on the fourth day his sedimentation rate was elevated to 30 mm and his electrocardiogram showed inverted T waves in all leads without the presence of Q waves in any lead. The T waves gradually became upright and in four weeks the electrocardiogram was again normal. He remained well during the next three years and his ECG remained unchanged after exercise.

A J, a sixty-three-year-old automobile salesman with a history of bronchial asthma for many years complained of sudden severe pain across the upper back radiating anteriorly. He perspired freely, his blood pressure was 130/80, a drop of 30 points from his usual systolic pressure. The pain lasted forty-five minutes and was relieved by 100 mg of Demerol subcutaneously. Despite a normal ECG he was suspected of having suffered an acute coronary occlusion. After four days he was permitted out of bed in a chair but he again began to experience frequent severe pain in the upper back during the day and particularly at night. Dicumarol therapy was begun and the patient received oxygen by mask, aminophyllin suppositories, Demerol 100 mg q 4 h and sedatives. The pain increased in severity and intensity but the patient's temperature, white count and sedimentation rate remained normal. On the eleventh day the ECG showed slight inversion of the T wave in lead I and V_{1-3} but no other changes. The T wave inversion gradually increased during the following two weeks and then receded four weeks after the acute episode the ECG had returned to normal.

In these cases coronary insufficiency with subendocardial ischemia without actual coronary occlusion clinically




FIG. 8-P I m 56 Acute coronary insufficiency in duodenal ulcer with profuse hemorrhage. Necropsy showed subendocardial necrosis. *A* Marked RS-T depression in all leads. *B* RS-T depression less marked. *C* Further improvement (Master Dick Horn, Freedman and Field, courtesy of Circulation.)

resembled a coronary thrombosis and in the second case anticoagulant therapy was started before the actual diagnosis of coronary insufficiency was made

In the more severe cases of coronary insufficiency there may be mild or moderate fever, leucocytosis and alteration of the sedimentation rate and longer lasting ECG changes. These may occur following operative procedures, massive hemorrhage¹⁷⁵ or tachycardias. Coronary insufficiency complicates gastrointestinal hemorrhage in many cases, especially when there is a marked fall in blood pressure (Fig. 8). For this reason we do not hesitate to give large amounts of blood to a patient who is actively bleeding, especially if he has coronary artery disease. All attempts should be made in these cases to keep the blood volume and blood pressure as close to normal levels as possible. Frequent cardiograms should be taken if acute coronary insufficiency occurs despite all efforts to prevent it; the patient should be kept in bed until the cardiographic abnormalities disappear. These cases usually recover, in the fatal cases necrosis of the subendocardial layer with evidence of coronary sclerosis is found, but there is no evidence of an acute thrombosis.⁸⁷

It is important to distinguish coronary insufficiency from coronary occlusion; in the former anticoagulant therapy is not usually necessary, the period of bed rest required is shorter and the prognosis is usually much better. In cases secondary to postoperative shock, massive hemorrhage, severe diarrhea or dehydration, treatment of the primary condition usually results in complete recovery from the cardiac difficulty.

Efforts to prevent attacks of severe coronary insufficiency should be made in all patients with any evidence of coronary disease. They should be advised not to indulge in strenuous activities; they should also be warned against excessive eating, exposure to extreme change in temperature, especially cold, frequent sexual intercourse, emotional upsets and excessive smoking. All attempts to prevent a fall in blood

pressure should be made especially when such patients are subjected to surgery. Spinal anesthesia may be used but care should be taken to prevent the blood pressure from falling below 100 mm Hg. If a rapid fall in blood pressure does occur during surgery due to the anesthesia or to massive hemorrhage blood should be given freely. If heart failure is induced by excessive fluid it may be quickly treated by digitalization the use of mercurials and the other measures previously described. It is far better to risk the occurrence of congestive heart failure which can be treated later than to allow a patient to lapse into irreversible shock by withholding transfusions merely because the patient has heart disease.

In diabetic patients with coronary artery disease who are receiving insulin it is important to prevent episodes of hypoglycemia. It may be better to allow them to spill small amounts of sugar than to attempt to attain perfect sugar control since hypoglycemic reactions often precipitate episodes of coronary insufficiency. If the above precautions are followed many cases of this syndrome will be prevented.

Coronary Occlusion —The pain experienced in coronary occlusion usually begins suddenly at rest during sleep or in the course of routine activity. It is usually precordial or substernal but may be situated only in the back of the chest or neck or in the arms. Instead of pain the patient may complain of severe burning sensations or feelings of tightness nausea vomiting and profuse perspiration. Varying degrees of shock may be present. The sudden onset of paroxysmal tachycardia or pulmonary edema is occasionally the first evidence of an acute coronary occlusion. Examination may reveal gallop rhythm embryocardia (tic tac rhythm) or congestion of the lungs. The blood pressure may be normal at first or it may drop precipitously.

The clinical and electrocardiographic manifestations of coronary occlusion depend upon the pathological changes that occur in the myocardium. In over half the cases the




FIG 9 —\ II m 45 RS-T depression and T wave inversion in the chest leads indicating coronary insufficiency during the premonitory phase of coronary occlusion (5 18 21) Following the acute attack RS-T elevation and Q waves appeared in leads V₂ and V₄ (5 22) indicating anterior infarction (Jaffe Courtesy of J Mt Sinai Hosp)

process of thrombosis is spontaneously initiated by a sub intimal hemorrhage which leads to secondary changes in the intima or may completely occlude the vessel lumen¹²⁰ This process of thrombus formation may take days or weeks During this period the patient may experience severe episodes of angina or recurrent pain at rest the so called premonitory phase of an occlusion¹²¹ (Fig 9) The electrocardiogram may show RS T depressions and T wave inversions indicating coronary insufficiency with myo




FIG 10 J S m 58 Acute coronary occlusion with postero lateral infarction (Master and Jaffe courtesy of Postgraduate Medicine)

cardial ischemia These changes are replaced by Q waves and ST segment elevations however as soon as the occlusion has become complete (Fig 10) A Q wave usually appears in one or more leads depending upon the site of the infarction and produces a characteristic pattern of Q and RS T elevation During the following days and weeks serial tracings reveal a progressive return of the RS T segment to normal and an inversion of the T wave resulting in a Q inverted T pattern which may persist for months or years Fever appears on the second or third day The sedimentation rate becomes prolonged by the fourth day, often to 50 mm or more In a through and through infarction the subpericardial region is involved and a pericardial

reaction²⁴⁶ and friction rub may result. Since the infarct reaches the intima mural thrombi may form and peripheral embolization occur.¹¹²

As we have mentioned previously it is oftentimes extremely difficult to differentiate between cases of simple coronary insufficiency and those which represent the premonitory phase of coronary occlusion. Both types therefore should be put to bed and observed before a diagnosis is made. In many cases of course where Q waves appear within eight to twenty-four hours following the onset of pain a definite diagnosis can be established quickly. It is also extremely difficult in many instances to distinguish between a coronary occlusion and an attack of acute pericarditis. Some of the features that differentiate these conditions from one another and from coronary insufficiency are summarized in Table II.

When electrocardiographic and clinical evidence of a coronary occlusion with myocardial infarction is present therapy should be begun immediately. Pain should be alleviated before other measures are instituted. If it is mild Demerol 50 to 100 mg or codeine 0.3 to 0.6 gm (gr $\frac{1}{2}$ to 1) should be given orally and will usually suffice. In more severe cases however morphine 15 mg (gr $\frac{1}{4}$) subcutaneously will be effective but it may occasionally be necessary to give morphine 10 to 15 mg (gr 1.6 to 1.4) intravenously. These doses may be repeated in twenty to thirty minutes. The possibility of inducing distention, obstipation and nausea and vomiting with morphine is great and the dose used should be no larger than is actually required. The morphine should be discontinued as soon as any of the above symptoms appear because distention and vomiting may turn an otherwise benign course into a stormy one. We have used Dilaudid extensively and especially in cases where morphine induced vomiting we have found it to be a reliable analgesic and to be without the unpleasant and dangerous side effects of morphine. Two and one half to 5 mg (1.24 to 1.12 gr) usually produce

quick relief of pain. We have had some experience with Pantopon and Metapon, opiate derivatives that produce few toxic reactions and have found these drugs also to be satisfactory. Nitroglycerin should not be used for the control of pain once a diagnosis of coronary occlusion has been made since it may further lower blood pressure and decrease coronary blood flow.

TABLE II

	PAIN	SEVERE SWEAT	GALLOP RHYTHM	PERICARDIAL RUB	HEART FAILURE	ARRHYTHMIAS	DIP. BLOOD PRESSURE
Coronary Insufficiency							
a) Mild Angina Pectoris	Few seconds to 10-15 minutes	Absent	Usually absent	Absent	Absent	Absent	Absent
b) More Severe—Spontaneous or Secondary	May be more prolonged	May be precipitating factor e.g. hemorrhage	May occur	Absent	Uncommon but may complicate due to tachycardia or hemorrhage	Uncommon	Moderate and prolonged in rhythm tachycardia
Coronary Occlusion	Usually severe longer than 15 minutes may rarely have no pain	Present in about 10 per cent of cases	Commonly occurs	Common	Common	Both atrial and ventricular arrhythmias may occur	Often marked
Forward type	Mild rarely severe	Rare	Rare	Usually present	Uncommon	Atrial flutter and fibrillation	Rare may occasionally tamponade occur

Oxygen should be used in all cases where the pain persists after morphine has been given. A tent should be employed where possible, especially when prolonged oxygen therapy is anticipated because the use of a mask or nasal catheter often increases the patient's anxiety. Occasionally, intravenous aminophyllin, 0.5 gm. (7.5 gr.) is of value. In rare instances, sodium amytal, 0.5 gm. (7.5 gr.) by vein, has been effective when all other measures have failed.

The patient should be put to bed in a quiet room if possible, and all distracting influences such as telephone and mail, should be banned. The importance of obtaining prolonged physical and mental rest should be explained to the patient. Visitors should be limited to members of the immediate family for the first two to four days.

The most serious complication of coronary occlusion is

TABLE II

Fever	Leukocytes	Elevated serum creatinine level	ECC R-T Fl R-T D p Q-Waves	Pathology	Precipitating factor	Treatment
Absent	Absent	Rare	R-T Fl Absent R-T D p Transitory Q-Waves Absent	No pathological changes	Frost on chest cold in general etc.	Active
Usually present	May be present 11000- 15000	Moderate	R-T Fl Rare R-T D p (common) Q-Waves Absent	Subendocardial necrosis	1. Tachycardia 2. Hemorrhage 3. Spontaneous	Treat hypertension
Usually present after 36 hours	Usually present after 36 hours may be over 15000	Usually after 36 hours	R-T Fl (common) R-T D p Reversible Q-Waves (common)	Through and through infarction of myocardium	Usually mild if severe poisoning	Anticoagulants treat
Common	Depending upon etiology If viral leucopenia present	Common	R-T Fl Common R-T D p Rare Q-Waves Absent	No myocardial damage	Unknown rheumatic fever viral factor all traumatic	Treat primary disease

shock, which is due to a sudden failure of the heart to pump blood to the tissues, i.e., a fall in cardiac output and is not due to blood loss or to a decrease in venous return. This so-called 'cardiogenic shock' results from myocardial weakness caused by the infarct and from certain poorly understood reflexes. Theoretically, then, the use of transfusions or intravenous fluids is not indicated in these cases and in fact may lead to a further increase in venous pressure.

and congestive heart failure. Actually however the danger of inducing failure thereby is not great as has been shown by several recent studies^{72,73}. Although this danger has also not proven significant in our own experience failure only occurring in a small percentage of cases the administration of intravenous fluids has reversed the shock process and changed the eventual outcome in only a few cases. Intra arterial transfusions may prove more useful although sufficient data have not yet been accumulated with this method to judge its value. The patient in shock should be placed in a moderate Trendelenburg position and should be given oxygen. If he complains of severe pain morphine Dilaudid or Pantopon should be given as required. Small doses of neosynephrine 3 to 5 mg. may be given intramuscularly and may be repeated every two to three hours. This drug raises blood pressure without affecting the myocardium adversely and is therefore safer than epinephrine. If neosynephrine is not available Paredrine 5 to 10 mg (1/12 to 1/6 gr.) or norepinephrine may be used. If a patient remains in shock despite these measures for more than an hour or two or if only minimal improvement occurs small transfusions should be given. If blood is not available plasma or hypertonic glucose solutions should be administered until blood is obtained. These measures are employed in an effort to prevent irreversible shock and death although they are effective in only a small percentage of cases. One of us (A. M. M.) has used ACTH and cortisone in this condition in a few instances but the results thus far have been poor. Usually the patient will recover from the shock state within a few hours or will expire. Occasionally patients have remained in shock following a coronary occlusion for as long as 8 to 12 hours and have recovered but this is not the usual course of events. If pain is promptly relieved oxygen given and the other simple measures outlined above are carefully followed the percentage of patients who recover will be as high as can be expected from any form of treatment in this condition.

Clinical evidence of mild left ventricular failure is very commonly observed early in the course of coronary occlusion, a few râles are found in the chest and slight congestion of the lung fields is seen on x ray. As a rule, this type of failure requires no specific therapy or responds promptly to moderate salt restriction and to an injection of a mercurial diuretic. Occasionally, however, acute pulmonary edema is the first sign of a coronary occlusion or severe heart failure develops within several days after the occlusion. The treatment of heart failure in the course of coronary occlusion and in the course of chronic heart disease is the same, except for the use of digitalis, concerning this, there is a difference of opinion. Many physicians believe that digitalis should be administered routinely in congestive failure occurring during a coronary occlusion, just as if the latter were not present.⁸⁶ There are reasons, however, for exercising caution in the use of this drug: the myocardium in the region of the infarct is more irritable, and digitalis may increase the irritability inducing premature beats, ventricular tachycardia, and even ventricular fibrillation. Many of the episodes of ventricular tachycardia in coronary occlusion have occurred in patients receiving digitalis and the association is probably not entirely coincidental. In the course of a coronary occlusion, therefore, we first employ other measures for combatting the associated congestive failure. In pulmonary edema we give oxygen, morphine and, if necessary, intravenous aminophyllin and a mercurial diuretic. If relief is not obtained, however, we do not hesitate to inject Strophanthin K (or Ouabain, Strophanthin G) intravenously, the initial dose being 0.25 mg. This dose may be repeated within one hour, but the drug should not be used if digitalis had been given within the previous forty-eight hours. In unusual instances, if it is felt that digitalization is absolutely necessary, doses of 1 mg. may be given even if there is a history of recent digitalis intake. In less acute failure we employ digitalis only if mercurial diuretics and a salt poor diet do not produce improvement. In acute

coronary occlusion if we must digitalize the patient we do it somewhat more slowly than usually but otherwise follow the plan outlined in Chapter 1 p 21

Paroxysmal arrhythmias of all types frequently occur in coronary occlusion. Premature beats auricular fibrillation and flutter and supraventricular tachycardia are most commonly encountered. They usually do not alter the prognosis because as a rule they are of brief duration and remit spontaneously. For this reason it is not necessary to begin treatment of these arrhythmias for several hours except in patients with heart failure or shock.

M W a retired business man of seventy nine developed a coronary occlusion with anterior infarction and did well for two days. On the third day his heart rate suddenly increased to 180. The LCG showed a supraventricular tachycardia with bundle branch block. The arrhythmia was remittent for several hours and then was constant for four hours the patient became quite uncomfortable and dyspneic. It was decided therefore to administer intramuscular quinidine 0.4 gm (6 grains) but by the time this had been drawn into the syringe the arrhythmia had ceased and did not recur.

Treatment was withheld in this patient whose tachycardia remitted spontaneously. In cases however in which the tachycardia persists all available measures should be employed to stop it. Also in all cases with heart failure and shock treatment of the tachycardia should be instituted immediately. The arrhythmias are treated as described in Chapter 1. Sedatives should be used liberally in these patients because the anxiety resulting from the rapid heart rate superimposed on the recent severe chest pain may prove difficult to manage.

Ventricular tachycardia in acute coronary occlusion responds to Pronestyl and quinidine and these drugs should be given as outlined above (Chapter 1 p 35). A particularly difficult case was recently treated successfully with Pronestyl.

M G a sixty year old tailor was admitted to the hospital because of melena and hematemesis of several days duration caused by a duodenal ulcer. For several months he had experienced pain in the left arm on effort this symptom increased following the onset of bleeding. Physical examination revealed a blood pressure of 114/65 and was negative. The electrocardiogram however showed the pattern of an acute coronary occlusion with an anterior wall infarction and regular sinus rhythm. The course of the disease was uneventful until the twelfth day when the patient became dyspneic and perspired profusely. His heart rate was 190 and the rhythm was slightly irregular. The electrocardiogram showed a ventricular tachycardia. The patient received morphine intravenous quinidine 0.65 gm (10 grains) on two occasions intravenous magnesium sulfate and intramuscular quinine without effect. He developed congestive failure within twenty four hours and was given Mercuhydrin and Digoxin 1.5 mg by mouth with some improvement in the failure. He then received quinidine 0.4 gm (6 grains) every 2 hours by mouth. The rate of the ventricular tachycardia ranged between 130 and 140. On the third day of the arrhythmia he was given 1 gm (15 grains) of procaine amide (Pronestyl) intravenously in fifteen minutes. During the injection a few sinus beats appeared but the tachycardia recurred a short time after the drug was discontinued. The next day 2½ gm of Pronestyl were injected intravenously in thirty five minutes and numerous sinus beats appeared. Within two hours regular sinus rhythm appeared and the patient made an uneventful recovery.

Many years ago it was suggested¹⁵⁴ that quinidine be routinely administered early in coronary occlusion to prevent ventricular tachycardia. We do not subscribe to this view because in our experience with 1000 cases of coronary occlusion who did not receive digitalis ventricular tachycardia occurred in only 0.5 per cent.¹⁷⁴ It therefore is not wise to employ a drug with potential toxic effects such as quinidine in an attempt to prevent a complication which is so infre-

quent. Similarly we do not administer quinidine for premature beats unless they are numerous or arise in several foci. Now that Pronestyl, a relatively harmless but effective drug, is available, it may be given as soon as a significant number of ventricular premature beats appear, but it too should not be used prophylactically in all cases of coronary occlusion to avoid ventricular tachycardia.

Varying degrees of P-R interval prolongation and evidence of partial or complete heart block may occur²⁵ following a coronary occlusion, but these findings usually disappear spontaneously and are of no clinical importance.

Anticoagulant Therapy — The major advance in the treatment of coronary occlusion during the past decade has been the introduction of anticoagulant therapy to prevent pulmonary and peripheral emboli. The frequency of this complication in coronary occlusion has been reported to vary between 4½ per cent² and 60 per cent⁹ or more in pathological material, although death may have occurred from other causes in many of these cases. Anticoagulant therapy has been shown to reduce the frequency of embolic phenomena and of death in coronary occlusion by about 40 to 50 per cent.² It is our distinct impression that many severe cases of coronary occlusion, particularly those with congestive failure who have survived, would have succumbed without the aid of anticoagulant therapy, and we believe that this therapy should be used routinely in coronary occlusion if adequate laboratory facilities for performing prothrombin determinations are available.

Pulmonary emboli most frequently originate in the lower extremities and lodge in the lower lobes.⁶⁶ It is possible that a small percentage arise in the right auricle in the presence of auricular fibrillation or prolonged congestive failure. The symptoms of pulmonary embolism are very variable. There may be severe chest pain, dyspnea and collapse simulating coronary occlusion, or merely cough, fever or tachycardia (Table III). Electrocardiographic abnormalities are present in half the cases; the

TABLE III

	PAIN	DYSPNOEA	SEVERE SHOCK	HEMORRHOIDS	GALLOP RHYTHM	PERICARDIAL RUB	FEVER	ECG	X RAY
FULM MARY 1-month	often absent typical	often severe	common in cases with large infarct	present in 1 of cases	usually absent	absent	may be high	a) RS-T depression and T wave inversion b) coronary insufficiency c) Q3 positive	Many patches in lower lobe may be present
LOW MARY (Section V)	usually severe subcostal pre- cordial 1 ft arm	often mild rarely absent	occurs in 5-10% of cases	absent unless pulmonary embolism present	common	may be present in about 50% of cases	usually not above 101° 10% Serial changes	RS-T elevation & Q waves Serial changes	may show minimal consolidation or be completely normal



166 11—E V 5 40 As the pulmonary embolism fifteen days following pelvic operation Death nine hours later Necropsy revealed bilateral pulmonary embolism of right main and left lower lobe branches The posterior papillary muscle of the left ventricle showed focal necrosis 1 M—Record taken soon after onset showed RS-T depressions 2 M—Record several hours later showed prominent S wave in leads I and II and RS-T rounding in lead III (Dack Master Horn Gishman and Field Courtesy of Am J Med)

commonest single pattern being that of coronary insufficiency : *i.e.* RS T depression and T wave inversion²⁴ (Fig 11) Other patterns are those of acute cor pulmonale : *i.e.* a large S₁ and Q₁ with RT₂ elevation and occasionally of right bundle branch block Pulmonary emboli usually do not occur until after the first week of the attack most commonly during the second week About one third of the patients with pulmonary embolism expire within a few minutes or hours despite all treatment but if a patient survives for more than three to four hours he probably will recover²⁵ In many cases it is extremely difficult to distinguish between pulmonary embolization and fresh myocardial infarction The following points will be helpful in making a differential diagnosis

Although fever tachycardia and chest pain may be present in both conditions and cough may occasionally occur in cases with coronary occlusion and mild congestive heart failure the presence of cough and hemoptysis favors the diagnosis of pulmonary embolism Calf tenderness or a positive Homan's sign¹ indicates the presence of a thrombotic process in a lower extremity and therefore possible pulmonary embolization A loud pulmonic second sound not previously heard may indicate pulmonary hypertension secondary to infarction and the findings of a hazy patch of infiltration on chest x ray may further suggest the diagnosis of an embolic episode Electrocardiographic changes as described above may be present in pulmonary infarction but characteristic Q waves are absent All of these findings may help to distinguish between the two conditions Occasionally a pulmonary infarction occurs during the course of a coronary occlusion without the appearance of chest pain cough and electrocardiographic findings In these instances a persistently high fever or persistent leucocytosis may suggest the diagnosis

Arterial emboli to the brain the abdominal viscera and the peripheral vessels arise in mural thrombi in the left ventricle at the site of the infarction Since these thrombi

probably begin to form very soon after the onset of the attack anticoagulant therapy should be started as early as possible

The two anticoagulant drugs commonly used are *heparin* and *Dicumarol*. The former is given by injection and its effect is evident within an hour or two¹⁰. *Dicumarol* is administered orally produces a definite effect in thirty six to forty-eight hours and a maximum effect in seventy two to ninety six hours. We believe that its action begins in twenty four hours in many cases although the ordinary laboratory tests may not detect this early effect. *Heparin* alone may be used throughout the attack but because its injection is often painful and because it is expensive most physicians give both *heparin* and *Dicumarol* during the first two days of the attack and thereafter maintain the patient on *Dicumarol*. In this way an immediate anticoagulant effect is produced by the *heparin* which after the third day is maintained by the *Dicumarol*.

As soon as a diagnosis of coronary occlusion is made the patient receives 100 mg of *heparin* subcutaneously or 400 mg of *Depo heparin* intramuscularly as well as 300 mg of *Dicumarol* orally. A control prothrombin time is taken. When given subcutaneously the dose of *heparin* is repeated every eight hours but when the intramuscular preparation is used a second dose need not be given for twenty four hours. In many cases we have not employed more than one dose of intramuscular *heparin* and it is our impression that the results obtained are just as satisfactory as when repeated doses are given. There is usually no need to use intravenous *heparin*. In the above dosage *heparin* usually prolongs the clotting time to approximately double the control value i.e. from fifteen to twenty five minutes (Lee White method) and there is ordinarily no danger of hemorrhage. The clotting time should be determined immediately before in order to obtain a pre treatment level and two hours after the first dose of *heparin* in order to discover patients who are sensitive to it. Thereafter this determina

tion is unnecessary. If hemorrhage does occur in the course of heparinization a 1 to 2 per cent solution of protamine sulfate is given by vein at a dose of 1 mg of protamine per mg of heparin being administered. After the initial dose of 300 mg of Dicumarol a dose of 200 mg is given on the second day. The administration of Dicumarol requires much closer supervision than does heparin since there is a tremendous variation in the response to it from person to person and in the same person from day to day. Even after some stabilization of dosage appears to have been achieved further administration of the drug produces unpredictable effects in some cases. Therefore during the first two weeks of treatment the prothrombin time should be determined daily before the next dose of Dicumarol is given. In the majority of patients the dosage of 300 mg followed by 200 mg is quite safe but occasionally prolongation of the prothrombin time to fifty or more seconds occurs within three or four days and profuse hemorrhage results. For example

L. W. a woman of twenty eight with chronic rheumatic valvular disease mitral stenosis congestive failure and auricular fibrillation experienced many episodes of peripheral embolization. Following a recent one she received 300 mg of Dicumarol on the first day and 200 mg on the second. Her prothrombin time was thirty six seconds on the third day and no Dicumarol was given on that day or on the next. On the fourth day diffuse ecchymoses appeared over the entire body and her prothrombin time was ninety four seconds. She received 2 doses of 75 mg of intravenous vitamin K within five hours and the prothrombin time fell to normal limits within eight hours. An uneventful recovery followed.

M. R. a garage owner of fifty six developed a coronary occlusion. He received 300 mg of Dicumarol on the first day and 200 mg on the second. His prothrombin time rose to fifty two seconds and did not fall during the following two weeks although he received no more of the drug.

These cases illustrate the extreme variability of response that may occur following Dicumarol therapy.

In Dicumarol therapy the prothrombin time should be maintained at about 2 to 2½ times the control time i.e. between twenty five and thirty five seconds. In most patients this result is achieved with a daily dose of 50 to 150 mg. Urinalyses should be done every two to three days on patients receiving Dicumarol. If a significant number of red cells are found in the urine the dosage should be reduced or the drug temporarily stopped. If the prothrombin time is prolonged to sixty seconds or more or if bleeding occurs the drug should also be stopped and the patient given intravenous vitamin K. Hykinone (vitamin K) 72 mg. is administered intravenously and this dose is repeated in four hours if indicated. Usually the prothrombin time will be reduced towards normal levels within four to eight hours after 75 to 150 mg. of vitamin K have been given. In more severe cases frequent transfusions of fresh blood are invaluable. Recently the use of vitamin K oxide has been advocated by some observers who have found it to be more efficacious than ordinary vitamin K preparations¹⁸⁹ in the prevention of serious hemorrhage.

Because of the danger of hemorrhage Dicumarol or any other anticoagulant substance should not be administered to patients with a history of hepatic, gastric or renal disease or with a hemorrhagic tendency.

Occasionally a patient will develop hemorrhage while on anticoagulant therapy despite the fact that his prothrombin time has been reported to be within safe limits. A good example of this is seen in the case of a fifty eight year old hotel proprietor who developed a coronary occlusion with anterior infarction and was given 300 mg. and 200 mg. of Dicumarol on the first two days. On the fourth day his prothrombin time was twenty seconds but he passed a large tarry stool and his hemoglobin fell to 8.6 gm. He was given a small transfusion slowly and recovered quickly.

Anticoagulant therapy should be maintained throughout

the period of bed rest and until the patient has become ambulatory. In no instance should Dicumarol therapy be used in an institution that is not equipped to do reliable prothrombin determinations. Despite the reliability of a laboratory, however, mistakes are occasionally made in determining the prothrombin time and in all cases the clinical picture should be considered more important than the chemical findings when treatment is ordered. It should be remembered that prothrombin times are not accurate when performed on patients who are receiving heparin in addition to Dicumarol. Although heparin primarily effects clotting time, it may significantly prolong the prothrombin time. For this reason and because a Dicumarol effect is usually not noted for 48 hours, it is useless to do prothrombin determinations during the first two days of treatment when the plan of therapy outlined above is followed. Some physicians continue anticoagulant therapy indefinitely in order to prevent recurrent coronary thromboses.⁸ We have had no experience with this type of therapy, but we do not believe that it has proven to be of value thus far.

Anticoagulant therapy has thus far been employed only in large medical centers, hospitals in the larger cities, and well equipped medical clinics, facilities for prothrombin time determinations being limited or unavailable in outlying rural areas and in small hospitals. Recent investigations of a bedside method of doing these determinations have demonstrated, however, that a simple capillary method utilizing rabbit brain thromboplastin yields results which, although not so accurate as those obtained by more elaborate techniques, are accurate enough for all practical purposes.¹⁶⁰ Approximately 700 patients have now been followed with this method and results appear to indicate that the procedure is feasible and useful as a rough guide in determining Dicumarol dosage. If further studies confirm the validity of this method, anticoagulant therapy need no longer be denied a patient because of expense, inconvenience, or lack of laboratory facilities.

Because of the disadvantages of heparin and particularly of Dicumarol several new anticoagulant drugs have been introduced and are receiving a clinical trial. *Paritol*²⁷⁶ a polysulfuric acid ester of polyanhydro mannuromic acid is used intravenously and has been shown to prolong the clotting time for from eight to twelve hours. Its anticoagulant effect persists 2 to 3 times as long as does that of heparin and it is only necessary to give one injection every eight to twelve hours. The dosage is about 7 times that of heparin. It should not be administered to patients with renal disease and there have been occasional mild reactions to the drug.

Tromexan^{39 41 111} an ethyl ether of Dicumarol acts much more quickly than does Dicumarol and is not cumulative. Its effect is apparent in eighteen to twenty four hours and when used with heparin the latter drug may be stopped after the first day. The dose used is about 6 times that of Dicumarol. We have found it difficult to maintain adequate prolongation of the prothrombin time with tromexan but experience with this drug is not yet sufficient for a complete evaluation of its usefulness.

The careful use of anticoagulant therapy has greatly decreased the incidence of embolization in our patients.

We have also decreased the incidence of embolization by permitting patients with myocardial infarction to move about more freely. While in bed the patients are advised to move their legs frequently if using a bed pan is difficult they are permitted to use a bedside commode after the second day if their condition is satisfactory. They are cautioned never to bear down or strain and to prevent distention laxatives (mineral oil cascara or Milk of Magnesia) are given freely after the second day following the occlusion. If a bowel movement does not occur after three days a low enema is given. In some instances it may be necessary to administer laxatives as well as enemas every two to three days during the first two weeks of bed rest.

If after three weeks the electrocardiogram has stabilized

the white count has become normal the patient is afebrile and no complications have occurred the patient is gradually permitted to walk. We do not depend upon the sedimentation rate to determine the total period of bed rest since frequently it does not return to normal for two or three or even twelve months *i.e.* long after the patients are ambulatory and have returned to work. We feel that early ambulation has also greatly contributed toward lowering the incidence of embolization.

If embolization occurs despite these measures the anti-coagulants are continued. In cases of cerebral embolization caution must be observed in administering these drugs. If a spinal tap does not reveal hemorrhagic fluid and if it is feared that further emboli will occur further anticoagulant therapy is justified. Sedation should be avoided in these patients and caffeine sodium benzoate $7\frac{1}{2}$ gr (0.5 gm) intramuscularly every 4 hours and aminophyllin 3 gr (0.2 gm) intramuscularly every 4 hours may be given. Other supportive measures should be employed oxygen should be used and parenteral feedings continued until the patient is able to feed himself.

When a peripheral embolism occurs intra arterial papaverine 50 to 100 mg every two to four hours given proximal to the occlusion is sometimes of value in increasing circulation in the affected extremity. We have recently obtained excellent results with intra arterial Priscoline 50 mg (2 cc) every four to six hours. Various ganglionic blocking agents also produce a high degree of vasodilatation and reduce vasospasm when administered intravenously. We have found that the Methonium derivatives C 5 (bis-trimethyl ammonium pentane dibromide) and C 6 (bis-trimethyl ammonium hexane dibromide) produce a longer lasting blockade and are less toxic than Etamon (tetraethyl ammonium chloride). C 5 (10 mg cc) and C 6 (25 mg cc) may be administered in 2-3 cc doses intravenously every 4 hours to maintain an adequate blockade. If these drugs are not available Etamon (100-300 mg) may be given every 4 hours or repeated sympathetic nerve blocks

employed. Although we have not as yet used intravenous Dibenamine in this condition this drug should be of great value as its adrenergic blocking effect usually lasts from 36 to 72 hours after a single dose of 5 to 7 mg/kg. In view of this fact Dibenamine will probably prove to be superior to other medications presently in use. If the patient is able to tolerate oral therapy the measures outlined above may be supplemented by the use of Priscoline 50 to 75 mg every 4 hours. 688A (N,N-Dibenzyl letachlorethylamine hydrochloride) a new Dibenamine derivative which is an orally effective sympathetic blocking agent⁶⁷ has not as yet been tried clinically in this condition but this should be done in the future. If the patient is not seen early and if gangrene has already occurred the leg should be packed in ice in preparation for amputation.

Embolectomy is the treatment of choice if it can be performed within eight to twelve hours after the acute episode. In expert hands results have been good and even massive saddle emboli lodged at the bifurcation of the aorta have been removed successfully.³⁸ It is a serious operative procedure particularly in patients with recent myocardial infarction but it is justified unless the patient is critically ill. A patient recently observed illustrates the fact that embolectomy can be performed successfully even in the presence of serious heart disease.

A sixty three year old tailor developed an acute coronary occlusion with anterior infarction. He did not receive Dicumarol. On the fifth day he complained of severe pain in the right leg which became mottled and cold up to the mid thigh. The femoral pulse was not palpable. Five hours after the onset of symptoms operation was performed by Dr. Gabriel Seley of New York and an embolus was removed from the femoral artery. Complete function of the extremity was restored and the patient made an uneventful recovery.

If embolectomy is not feasible and amputation becomes necessary it should be performed under local or block anesthesia after demarcation has occurred.

Chapter

4

SYNCOPE

SYNCOPE occurs in many conditions, and usually in patients without organic heart disease (Table IV). Two main mechanisms account for the temporary cerebral anoxia, which is the physiological basis of syncope: (1) a sudden fall in cardiac output, *e g.* following severe hemorrhage³⁶ (2) a marked decrease in peripheral resistance and a fall in blood pressure with a fairly well maintained cardiac output, *e g.*, neurogenic fainting¹⁰³. Neurogenic or vaso-depressor syncope, by far the most common, is produced by reflex peripheral vasodilatation, and may be precipitated by the sight of blood or by fright, pain or severe emotional disturbances⁷⁰. It is more frequent in men than in women. Premonitory symptoms such as nausea, sweating or lightheadedness usually precede the attack which occurs only if the patient is standing. This type of syncope can usually be terminated by laying the patient down with his head lowered. If the syncope lasts for more than ten to fifteen seconds clonic twitching may appear but true grand mal seizures are extremely rare. In this type of attack it is sometimes necessary to administer sympathomimetic drugs, *i e.*, epinephrine, 0.5 cc. of 1:1000 solution subcutaneously.

Another form of syncope due to peripheral vasodilatation is observed in patients who while sitting or recumbent employ heating pads for low back pain. When these patients suddenly resume the erect position a temporary cerebral anoxia occurs and fainting may result. Fainting

TABLE IV - DIFFERENTIAL DIAGNOSIS AND TREATMENT OF SYNCOPE

TYPE	PRECIPITATING FACTOR	PREMONITORY SYMPTOMS	DURATION OF ATTACK CONVULSIONS?	LABORATORY FINDINGS	TREATMENT
1) Vascular pressure response a) No organic	Fright, pain, emotional upset, sight of blood	Nausea, sweating, light-headedness, dizziness (only occurs when patient is standing); premonitory phase 1-5 minutes	Usually of short duration 1-15 seconds—after 15 seconds twitching may appear	Cardiac output normal Blood pressure falls Pulse rate may rise or fall Electroencephalogram abnormal	Usually terminated by laying patient down with head lower than feet Rarely 5 to 1 cc. Epinephrine (subcutaneously) if necessary
b) Carotid sinus syncope	a) Pressure on neck b) Turning head c) Cerebral arteriovenous aneurysm d) Distal atherosclerosis	Usually no premonitory symptoms	Short duration Convulsions may occur	Blood pressure falls Pulse rate decreases Electroencephalogram abnormal	Epinephrine Sulfate 25 mg. t.i.d. Atropine 3 mg. t.i.d. Excess of vagus nerves if attacks are severe and persistent
c) Secondary to Other Factors	Internal or external hemorrhage, fever	Same as (a)	Same as (a)	Same as (a)	Treat primary condition
2) Postural Hypotension	a) Talipes Equinovarus b) Spinal Cord Tumors c) Addison's Disease d) Diabetes Mellitus e) Standing in one place, arising suddenly f) Lethargic	Light-headedness, dizziness	Usually of short duration (1-5 seconds) No convulsions	Blood pressure falls Pulse rate increases Electroencephalogram abnormal	Lay the patient down Tight abdominal belt and neosynephrine or epinephrine Tilt bed Cautious against raising quickly Treat primary disease
3) Hysterical Syncope	a) Suggestion b) pressure on any area of body c) tense situation d) Hyperventilation	No premonitory symptoms	Never any convulsions May last for hours	No change in blood pressure or pulse rate No electroencephalogram changes	Psychotherapy
4) Cardiac Syncope	a) Heart block with periods of asystole b) Ventricular tachycardia or other arrhythmias	Diarrhea, palpitations, chest pain, nausea, epigastric pain	Usually short (5-10 seconds) May last longer with convulsions	Electrocardiographic abnormalities Blood pressure falls Pulse rate may be extremely slow or imperceptible	Treat card. arrhythmias. Quinidine, Digitalis, Proserin. For heart block: Epinephrine, Atropine etc. See text
Hypoglycemic Syncope	a) Islet cell tumor b) Overactive parathyroids c) Too great an interval between meals d) Hepatic disease	Weakness, extreme hunger, tiredness, sweating, pallor	May be prolonged if not treated Convulsions occur frequently	Blood sugar extremely low Electroencephalogram abnormal Pulse rate increased Blood pressure unchanged or low	Intravenous glucose then oral glucose—depending upon etiology (see text) a) remove tumor if present b) low carbohydrate diet c) 1 g. carbohydrate diet if liver disease is present
Tumors	a) Severe exophthalmos b) Breathing down	Diarrhea, tightness	Short (1-5 seconds)	No unusual findings	Warn patient against leaning down or exerting effort a) remove tumor

occasionally occurs in normal persons who are required to stand in one place for long periods of time and in patients with autonomic nervous system disturbances *i.e.* over active vagal reflexes (carotid sinus syncope) or poor sympathetic nervous regulation of blood vessels^{23 24} (postural hypotension)

Many persons with a hypersensitive carotid sinus faint when the sinus is massaged²³ but only a small percentage of these experience spontaneous syncope as a result of this sensitivity. Consequently a diagnosis of carotid sinus syncope should be made only if the following criteria are satisfied (1) The attacks must occur without the premonitory signs usually seen in the type of vasodepressor syncope not associated with carotid sinus hypersensitivity (*i.e.* syncope due to fright or sight of blood) (2) The attacks should be reproducible by sinus stimulation (3) The attacks should not occur after the administration of adequate doses of atropine or ephedrine *i.e.* doses large enough to prevent bradycardia on manual carotid sinus stimulation. True cases of this syndrome are relatively uncommon when compared with other types of vasodepressor syncope.

An interesting illustration of carotid sinus sensitivity is furnished by a seventy one year old man who was admitted to the hospital because of four attacks of syncope during the preceding four months. Examination revealed a blowing systolic murmur at the aortic area transmitted into the neck. The blood pressure was 160/80. The peripheral pulses were not palpable. The electrocardiogram showed evidence of previous lateral wall infarction. In the hospital he experienced brief episodes of dizziness. It was thought that his syncope was the result of cerebral arteriosclerosis and aortic stenosis. Pressure on the right carotid sinus however produced asystole and a drop in blood pressure to 80/0. The patient became blanched, his hands twitched and he yawned. Stimulation of the left carotid sinus produced similar results but with less regularity. He was given oral

atropine 3 mg (gr 1/100) q i d and after several days carotid sinus pressure caused no change in the heart rate (Fig 12) The patient has been maintained satisfactorily on this medication It was extremely difficult to establish a diagnosis of *true carotid sinus syncope* in this case despite the fact that definite *hypersensitivity* of the sinus could be demonstrated The patient had cerebral arteriosclerosis and aortic stenosis which alone may have caused the syncope




FIG 12—J Q m 61 Arteriosclerotic heart disease. Continuous lead II. A Control sinus bradycardia rate 52 B After pre-re-

Since atropine definitely prevented the attacks however, the diagnosis of syncope secondary to carotid sinus sensitivity, was the most logical one. Patients who have extensive arteriosclerosis and are receiving digitalis are more susceptible to syncopal attacks and, in some instances stopping the digitalis will eliminate the attacks completely. Carotid sinus syncope may be induced by pressure on the sinus from any cause *e g* by an enlarged gland in the neck or by a sudden movement of the head.

Tight collars often serve as a trigger mechanism. J C, a thirty eight year old midwestern preacher was investigated because he had fainted several times while delivering his Sunday sermon. In eliciting his history it was discovered that during the week he worked in the field with his collar open on Sunday, however he wore a stiff collar to appear in church. Carotid sinus pressure repeatedly induced asystole and syncope in this patient and the attacks were eliminated when the patient stopped wearing tight collars.

Convulsions may occur as a result of excessive carotid sinus stimulation²⁰⁴ and caution must be observed when testing a patient for sensitivity especially an older person or one with arteriosclerosis. Pressure should never be applied for more than ten seconds and the patient should be recumbent when tested. Ephedrine sulphate (25 mg) Paredrine (60 mg 3 times a day) or atropin sulphate (0.3 mg grain 1/200), will prevent most attacks. Neosynephrine, in oral doses of 20 to 25 mg 3 times a day is occasionally successful when other medication fails. When it is necessary to use sympathomimetic drugs *i e*, ephedrine for long periods of time, it is wise to administer small doses of sedatives (barbiturates) simultaneously to prevent nervousness and insomnia. In a few cases excision of the sinus has given permanent relief.¹⁵⁰

Postural hypotension may occur in normal persons, in patients with neurocirculatory asthenia and in those with serious underlying central nervous system disease,⁶⁸ such as cord tumor or tabes dorsalis.²²⁷ We have also observed

extremely severe postural hypotension in patients with advanced diabetes mellitus Addison's disease and pheochromocytoma. When these people assume the upright position the normal compensatory splanchnic constriction does not occur the cardiac output is diminished and the blood pressure falls.

In patients with chronic postural hypotension and recurrent syncopal episodes therapy is extremely difficult. The primary disease should be sought and treated if possible *i.e.* Addison's disease varicose veins *etc.* If no primary condition is found atropine epinephrine neosynephrin or Paredrine should be tried in the doses cited. These drugs have not proven very successful however and are usually more effective when used in conjunction with a tight abdominal belt and/or elastic bandages around the lower extremities. It is occasionally temporarily helpful for these patients to sleep in a bed tilted 20 to 25 degrees from the horizontal with the head up.⁶⁴ Desoxycorticosterone acetate and sufficient salt to cause slight dependent edema sometimes reduce the frequency of attacks. The patients should be instructed to rise from bed slowly and to avoid taking drugs that cause peripheral vasodilation *i.e.* amyl nitrite or nitroglycerin.

Fainting may be hysterical in origin particularly in women. This type of syncope is not accompanied by premonitory symptoms and is not associated with any changes in pulse or blood pressure. These patients may fall to the ground and remain unconscious for a period of several seconds or hours but convulsions do not occur and injury rarely results. Electroencephalographic changes which are seen in most other forms of syncope that last more than a few seconds do not appear in this condition.⁷⁰ Fainting may be precipitated by suggestion or by manipulation of almost any part of the body. Therapy is mainly psychiatric.

In anxiety states syncope occasionally occurs as a result of hyperventilation. Such an attack may be reproduced by having the patient breathe deeply for one or two minutes.

Assurance and sedation are the only treatments of value in correcting this condition

A rare form of syncope is that which follows severe bouts of coughing or the Valsalva maneuver (a forced expiration with the glottis closed—the bearing down effort)¹³ A marked rise in intrathoracic pressure occurs and prevents adequate filling of the right atrium resulting in a marked fall in cardiac output and cerebral anoxia. Recognition of this type of syncope is important since most of these attacks may be prevented by warning the patient against excessive bearing down efforts such as straining at stool and against remaining upright during a spell of severe coughing.

Fainting which occurs early in the morning before breakfast or several hours after a meal may be the result of hypoglycemia. This type of syncope is seen in cases of hyperinsulinism caused by pancreatic adenomata in patients on a high carbohydrate diet with an overactive pancreas and in cases of marked hepatic disease.⁴⁵ The patient usually experiences weakness, dizziness, and extreme hunger before fainting; convulsions often occur if therapy is not instituted quickly. Intravenous or oral glucose should be given as soon as possible and recovery from the attack is usually rapid. The diagnosis is made from the history, the fasting blood sugar determination, and the glucose tolerance test. Extremely low blood sugar levels may be obtained during the course of the latter test, especially in the two, three, and four hour specimens, and fainting may occur before the test can be completed. These attacks may be prevented in several ways depending upon the etiology of the condition. If hypoglycemic attacks occur before breakfast and rarely during the day, and if liver damage is demonstrated, a high carbohydrate, high protein diet should be prescribed. If the attacks occur before breakfast and frequently during the day, and if they become progressively more severe and are not relieved when the patient is put on a low carbohydrate, high protein diet, then a pancreatic adenoma is probably present and surgical removal of the tumor is advisable.

If the attacks occur approximately two to three hours after meals, and if no evidence of liver disease or progression is found, then so called 'functional' hypoglycemia is present, and the patient should be put on a low carbohydrate, high protein diet with frequent feedings. Additional points of differentiation between these three major types of hypoglycemia, and suggestions for therapy may be found in a recent review.⁴³

In organic heart disease syncope occurs in complete heart block, *i.e.*, Stokes Adams attacks, and sometimes at the onset of paroxysmal tachycardia⁴⁴ (Chapter 1 p. 11) and following an acute coronary occlusion. In patients with calcific aortic stenosis secondary to rheumatic or arteriosclerotic heart disease fainting may also occur especially following exertion. The prognosis of syncope in organic heart disease is, of course, more serious than it is in non cardiac cases. Therapy in the former group is directed against the primary condition.

Chapter

5

RHEUMATIC HEART DISEASE

DURING the acute stage of rheumatic fever the occurrence of acute pericarditis congestive failure and arrhythmias frequently necessitate emergency treatment. In such cases early adequate treatment often reduces the morbidity and mortality rate. Previously only salicylate therapy and symptomatic treatment were given. The recent introduction of ACTH and Cortisone however has provided the physician with drugs that appear to be extremely useful and rapidly effective in the treatment of the acute rheumatic process.^{2 1 70 269} For this reason these drugs should be used whenever possible in cases of active rheumatic carditis that are seen early.

We have recently observed a critically ill patient in severe left and right heart failure with rheumatic pancarditis who did not respond to the usual therapy but who improved remarkably after receiving 25 mg of ACTH every six hours for six days. There was little doubt that death would otherwise have occurred. After six days the daily dose of ACTH was reduced from 100 to 60 mg and this was maintained for seven days. The daily dose was then further reduced to 40 mg. This treatment was continued for two more weeks and the patient recovered.

A course of treatment of from seven to ten days has been suggested to control the carditis⁶⁹ but we feel that ACTH should be continued for a period of at least two to three weeks as was done in the case cited. If this does not prove effective another course of therapy may be given. If

If the attacks occur approximately two to three hours after meals, and if no evidence of liver disease or progression is found, then so called 'functional' hypoglycemia is present, and the patient should be put on a low carbohydrate, high protein diet with frequent feedings. Additional points of differentiation between these three major types of hypoglycemia, and suggestions for therapy may be found in a recent review.⁴⁵

In organic heart disease, syncope occurs in complete heart block, *i.e.* Stokes Adams attacks and sometimes at the onset of paroxysmal tachycardia⁴⁷ (Chapter 1, p. 11) and following an acute coronary occlusion. In patients with calcific aortic stenosis secondary to rheumatic or arteriosclerotic heart disease fainting may also occur, especially following exertion. The prognosis of syncope in organic heart disease is, of course, more serious than it is in non cardiac cases. *Therapy in the former group is directed against the primary condition.*

alter the subsequent course of rheumatic heart disease, i.e. the development of valvular lesions, congestive heart failure etc. is questionable.

Should *congestive failure* or *arrhythmias* occur during the course of acute rheumatic fever they are treated as has already been outlined. Digitalis however should be used with caution in these cases. This drug is often ineffective in the acute phase of rheumatic carditis and may even be harmful; furthermore the tendency is to push the dosage to toxic levels in order to obtain an effect.³ It is therefore wiser to depend upon salt restriction and mercurial diuretics in congestive failure before digitalization is attempted. If failure persists however despite these measures digitalis may be used. Although some observers² have reported favorable results with digitalis in the presence of active carditis we are unable to confirm these observations. The drug is occasionally effective in cases with congestive heart failure and auricular fibrillation but its administration should be carefully controlled.

Acute pericarditis may be associated with severe precordial or epigastric pain which is relieved only by morphine. The pain usually develops within a few days after the onset of acute rheumatic fever. The heart sounds are distant, a friction rub may be heard and shock may occur. The electrocardiogram usually presents a typical pattern consisting of RS-T elevation in two or more leads without Q waves (Fig. 13). If an effusion accumulates the rub may disappear and the patient become more dyspneic and orthopneic. Tapping is rarely necessary in these cases however and has been performed only when a rapidly progressive increase in dyspnea and venous pressure occurs as a result of cardiac tamponade. Usually the fluid is resorbed spontaneously as the acute infection subsides on bed rest and drug therapy.

In the course of chronic rheumatic heart disease with mitral stenosis and auricular fibrillation auricular thrombi may form and in rare instances occlude the valve open




FIG. 13 - A P f 26 Acute rheumatic fever with pericarditis. RS-T elevations are present in all leads.

ing^{27 171 226} (ball valve thrombus), producing a serious emergency. Cyanosis appears at the tip of the nose, the lobes of the ears, and the fingers and toes and severe dyspnea, congestive heart failure, and shock usually develop. Death may occur suddenly⁶ or after a few hours or days. The treatment is supportive, including oxygen and the usual measures employed in congestive failure and shock. Morphine 15 mg ($\frac{1}{4}$ gr), may relieve the restlessness. Occasionally a change in position brings relief to the patient. We have seen the cyanosis disappear immediately when the patient sat up and leaned forward.

Pulmonary infarction and cerebral and peripheral arterial occlusion may also be caused by auricular thrombi that are released into the blood stream. One embolic episode is often followed by others. In such cases, chronic Dicumarolization is indicated. The dosage is adjusted while the patient is hospitalized, for the next month prothrombin times are done 3 times a week, thereafter, weekly determinations are sufficient. The results reported with this treatment appear to be satisfactory.^{106 218} We have observed three patients during the past eighteen months who had experienced repeated bouts of embolization, and who had consequently been continuously bedridden. They were hospitalized and treated with Dicumarol being maintained on the drug after discharge. They have now been followed for from nine to eleven months, and have not experienced a single embolic episode, a remarkable result in view of the previous course of their disease. Despite the fact that chronic Dicumarolization may markedly decrease the number of embolic episodes this procedure should not be carried out if adequate facilities for determining prothrombin times are not available. Chronic anticoagulant therapy with the drugs now in use is by no means a simple, safe procedure, and should be embarked upon with great caution.

In 1946, Dock⁶⁸ suggested left auricular appendectomy or ligation in patients with mitral stenosis in order to reduce

the high incidence of embolization since he believed that most thrombi originated in this appendage. Recently the auricular appendage has been tied off successfully in five cases^{12, 169} although the results were poor in the first three¹⁶. In view of the fact that thrombi may form in both appendages in cases of rheumatic heart disease¹³³ especially in those with congestive heart failure it is quite possible that bilateral auricular resection will prove to be the correct surgical procedure to prevent embolization. Since neither of these operations is a simple one follow up studies will be necessary to justify their use. The results thus far obtained suggest that the continual use of anticoagulants produces the same effect as auricular resection without its serious drawbacks for this reason we favor the former treatment in these cases.

In rheumatic heart disease embolization should suggest the presence of subacute bacterial endocarditis. The emboli are generally small and occlude small peripheral arterioles producing white centered petechiae. Every patient with rheumatic heart disease who runs an unexplained fever for more than one week should have blood cultures taken. Three blood cultures may be taken during the first day it is unnecessary to wait twenty four hours between cultures as is commonly done. If a patient with rheumatic heart disease exhibits white centered petechiae or splenomegaly in addition to the fever a diagnosis of subacute bacterial endocarditis is justified even if blood cultures are negative and therapy should be promptly instituted⁸¹. In a case recently observed by us cerebrovascular embolization occurred because treatment was delayed pending the report of the blood cultures. In any case with clinical findings of subacute bacterial endocarditis therapy should be started after two blood cultures have been taken i.e. eight hours after hospitalization.

Recently in patients with tight mitral stenosis several surgical procedures have been employed in an effort to decrease the incidence of intractable congestive failure.

hemoptysis and recurrent attacks of pulmonary edema. Such attacks are extremely frequent in patients with mitral valvular lesions, especially following activity.¹⁶ An extra-cardiac shunt to prevent pulmonary edema by means of an anastomosis between a branch of the inferior pulmonary vein and the azygos vein has been performed with some good results,²⁷ but this operation has now been replaced by direct operation on the mitral valve.

Successful operative repair of the mitral valve has been done¹¹⁰ in patients with mitral stenosis whose cardiac output was low and whose pulmonary venous pressure was markedly elevated, i.e., patients with a 'tight' mitral stenosis with little insufficiency. In cases with normal cardiac output, i.e., with a lesser degree of stenosis, it is sometimes better to create an interatrial septal defect. When patients are too sick to be operated on, repeated sympathetic blocks may be of value in preventing paroxysmal tachycardia and or acute pulmonary edema.¹¹⁰ Much greater experience is required before definite conclusions can be drawn regarding the operative correction of mitral stenosis.

Chapter

6

HYPERTENSIVE ENCEPHALOPATHY AND CRISES

HYPERTENSIVE crises may occur in the course of essential hypertension particularly in its malignant phase and in acute nephritis. The patient becomes acutely ill with unremitting headache and may exhibit temporary blindness paralysis aphasia and convulsions. These attacks are often brief but if they persist relief may occasionally be obtained by the intravenous administration of 20 cc of a 10 per cent solution (2 to 4 gms) of magnesium sulphate by withdrawal of spinal fluid or by the intravenous injection of 50 cc of 50 per cent dextrose. Good results have also been achieved with intravenous Dibenamine (5 to 7 mg per kg in 100 cc of 5 per cent glucose in water given in one hour) ^{191 275}. This drug lowers the blood pressure one dose may alleviate the mental symptoms for several days. The infusion of Dibenamine may be repeated every forty-eight to seventy two hours to keep the patient symptom free.

Pheochromocytoma a tumor which secretes both epinephrine and nor-epinephrine is usually found in the adrenal medulla or the retroperitoneal chromaffin tissue. It may cause attacks of severe hypertension with headaches palpitations tachycardia vertigo abdominal pain nausea vomiting and profuse sweating ¹⁹. Because of the increased amount of circulating adrenalin in patients with a pheochromocytoma an elevation of the basal metabolic rate sometimes occurs. Occasionally glycosuria and an elevated blood sugar are present as a result of an increased glycogen

breakdown secondary to epinephrine action. The attacks of palpitation sweating *etc* may be sudden in onset and occasionally can be induced by palpating the region of the kidneys. The patient becomes acutely ill and pale and breaks out in a cold sweat. In cases of pheochromocytoma the hypertension may be persistent and chronic headaches and continuous sweating may be present. The diagnosis of this tumor should be considered in hypertensive patients who perspire profusely or who have an elevated basal metabolic rate a glycosuria or a marked loss of weight. It may be confirmed by a positive response to histamine¹⁴⁸ or tetra-ethyl ammonium chloride (Etamon)¹⁴⁹ i.e. a rise in blood pressure and a reproduction of the acute attack. The rise in blood pressure is probably the result of a dilatation of blood vessels in the tumor and an increased outpouring of adrenalin. Benzodioxane⁹⁸ and Dibenamine¹⁴⁹ which cause a fall in blood pressure and bring relief of symptoms as a result of their adrenolytic action are also extensively used as aids in establishing the diagnosis. Benzodioxane at present appears to be the most satisfactory diagnostic agent during the hypertensive stage of the disease. A false positive reaction rarely occurs when it is used. If hypertension is not present when the patient is tested Etamon is preferable to histamine because it produces a less violent rise in blood pressure. Although calcification in this tumor is uncommon its presence in the region of the adrenal gland may suggest the true diagnosis in a hypertensive patient with the above mentioned symptoms.¹⁴⁷ Since hypertension secondary to a pheochromocytoma is the only type of hypertension that can be cured by surgical intervention we feel that *all* patients with either fixed or paroxysmal hypertension should be studied for the presence of this tumor. The testing agents now available are simple to use and do not necessitate hospitalization of the patient.

The acute attacks should be treated with intravenous Dibenamine given as described above. The tumor should be excised as soon as possible.

Chapter

7

DISSECTING ANEURYSM

DISSECTING aneurysm of the aorta is relatively uncommon but its diagnosis is important for two reasons (1) if an erroneous diagnosis of acute coronary occlusion or pulmonary infarction is made and anticoagulant therapy is given the possibility of recovery from a dissecting aneurysm is further reduced (2) if a method for the surgical treatment of dissecting aneurysms is perfected early diagnosis will be essential for success. Only a small percentage of the cases is diagnosed antemortem probably because of the failure to suspect the presence of the condition. Contrary to general belief a dissecting aneurysm is not invariably fatal 10 to 25 per cent of patients have lived for from three months to eight years following its onset ²⁵ The dissection may be acute lasting only a few hours and terminate in death as a result of rupture of the aorta or it may be subacute causing repeated bouts of pain for a period of 2 to 3 days and end in recovery.

The attack does not appear to be precipitated by unusual exertion usually occurs in patients with hypertension and is sudden in onset. Occlusion of a branch of the aorta *i.e.* iliac intercostal or coronary is common. There is a sharp knife like pain in the chest and/or back with occasional radiation to the epigastrium the lower abdomen the flanks and the legs. Syncope is noted in 10 per cent of the cases and shock may occur quickly. The patient may writhe in pain but unlike the pain of coronary occlusion this pain is not crushing or constrictive and does not

require a minute or two to reach its maximum intensity. Various neurologic signs may appear including paraplegia which is caused by hemorrhagic infarction of the spinal cord.²¹⁰ An aortic diastolic murmur is heard in many cases. Electrocardiographic evidence of left heart strain or coronary insufficiency may appear or of coronary occlusion if the dissection occludes a coronary artery. A coronary occlusion secondary to a dissecting aneurysm may sometimes be differentiated from the ordinary type of coronary occlusion by the presence of pain in the back, the lower abdomen or the legs, by the sudden appearance of a diastolic murmur or by the presence of neurologic changes. If a pleural or pericardial effusion or hemorrhage into the wall of the aortic arch occurs, a widened mediastinum may be detected on fluoroscopy or x ray. This procedure should be carried out with great caution and actually should not be used if the diagnosis is fairly well established from other findings. The diagnosis should be considered in any patient with hypertension who suddenly develops severe knife like pain in the chest, back, epigastric region or legs, and in whom a diastolic murmur suddenly appears.

A case in which the correct diagnosis was suggested during life was that of a hypertensive man of 42 who developed episodes of severe substernal pain and shock with apparent complete recovery between attacks. The electrocardiogram showed the deep Q waves and RS T elevations characteristic of coronary occlusion. On the third day, during one of his attacks, the heart sounds suddenly became faint and the area of cardiac dullness increased. A pericardial tap revealed hemopericardium and a diagnosis of dissecting aneurysm was made. He died almost immediately and postmortem examination disclosed a dissection involving a coronary artery.

Many times, however, the history and findings are completely atypical and a correct diagnosis cannot be made. For example, A sixty three year old white male complained of sudden, severe, intermittent pain in the left lower

quadrant. He had experienced mild pain in that region eight months before for a period of four or five days but the pain had not recurred until the present admission. Examination revealed an ill defined mass in the left lower quadrant. His blood pressure was 170/100 and no murmurs were heard. A diagnosis of carcinoma of the colon or diverticulitis was made. Several days later the patient was found to be quite pale and the mass appeared to be larger. Twenty four hours later he died suddenly. Necropsy showed evidence of dissection of the aorta from the thoracic to the abdominal portion with rupture of the abdominal aorta and retroperitoneal hemorrhage.

When the presence of a dissecting aneurysm is suspected the patient should be put to bed immediately. Repeated doses of morphine should be given if necessary to attain complete sedation. He should be kept in bed for at least three weeks and then should gradually be permitted to walk. Anticoagulants should not be used. If an accurate diagnosis is made and if treatment is instituted early a much larger percentage of cures may be effected.

In view of the tremendous advances in vascular surgery it is not unlikely that a procedure will be developed to rechannel the blood in the dissected coats of the aorta into the lumen of the vessel. A prerequisite to this however is the discovery of methods to detect the site and path of the dissection.

Chapter

8

TRAUMATIC HEART DISEASE

THE heart may be injured directly by penetrating bullet or knife wounds or indirectly by severe blows to the chest^{10 17}

The diagnosis of a penetrating wound of the heart is simple if external evidence of the injury is present. Occasionally, however, a bullet or a needle which had entered the body at some remote point, may be carried to the heart through the veins,¹⁰⁸ and produce cardiac damage. In these cases the diagnosis is often difficult. If a patient with a recent penetrating injury of the chest wall presents evidence of blood loss, congestive heart failure or shock, trauma of the heart should be suspected. The commonest result of such trauma is hemorrhage into the pericardial sac or laceration of the myocardium with infarction. In rare instances valvular rupture may occur. When a hemopericardium develops, fluoroscopy or x ray may show evidence of a widened cardiac shadow with diminished pulsations or electrocardiograms alone may show characteristic changes. Occasionally findings in such cases are similar to those seen in myocardial infarction. Surgical intervention is justified only when there is evidence of increasing hemopericardium and cardiac tamponade: i.e., shock, with a weak thready pulse and drop in blood pressure (low cardiac output), or congestive heart failure. A pericardial tap should be performed immediately, and as much blood as possible should be aspirated.²¹⁴ If blood reaccumulates, or if the symptoms

persist or recur a severe myocardial laceration has occurred and should be repaired immediately

Shock should be treated as efficiently as possible before and during the operation. The Trendelenburg position, frequent transfusions of blood (plasma if blood is not available) and the liberal use of morphine intramuscularly or intravenously will often help the patient to surmount the crisis until the wound is repaired. When plasma or blood is not immediately available intravenous infusions of 10 per cent glucose in water may be given in the meantime. Despite the presence of cardiac tamponade the cardiac output may be increased and the patient's condition improved by these measures.⁹

Excellent reviews of the technique of repairing lacerations or removing foreign bodies have been written.⁷⁻⁹ These surgical procedures need not be discussed here. Complications such as cardiac arrest encountered during operation are discussed in Chapter 9. In experienced hands the operative mortality rate in such cases is 25 to 30 per cent.⁶ The site repaired heals and a firm scar develops. Despite the fact that a major coronary artery is often ligated infarction does not always occur and cardiac function is usually good.¹⁷ A case seen by us demonstrated this finding. W. C., a thirty-five year old laborer was stabbed in the region of the heart with a long knife two hours prior to admission. There was profuse hemorrhage but the patient walked several blocks to the hospital. He complained of pain in the chest but was neither dyspneic nor cyanotic. The blood pressure was 100/80. The heart sounds were distant and a pericardial rub was heard which soon disappeared. An x-ray of the chest and an electrocardiogram were negative. In view of the profuse bleeding an exploratory operation was performed. A laceration of the parietal pericardium with a moderate hemopericardium and an insignificant stab wound of the anterior surface of the heart were found. Upon gentle manipulation the oc-

cluding blood clot was dislodged and free bleeding occurred this could be controlled only by the insertion of three sutures, two of which entered the chamber. Following the operation, the electrocardiogram showed RS-T elevation followed by T wave inversion in the standard leads (chest leads were not taken because of the chest strapping) the typical serial changes of acute pericarditis. In addition, a Q wave appeared in lead II this produced a W shaped QRS which was still present a year later when the T waves in the standard leads had returned to normal. It is possible that the alteration in the QRS was produced by the injury associated with the suturing. Despite the fact that a coronary artery had been ligated during the operation the patient made an excellent functional recovery.

Nonpenetrating chest injuries may produce signs and symptoms that are difficult to distinguish from those found in spontaneous coronary occlusion or coronary insufficiency. The causal relationship between trauma and myocardial damage is frequently difficult to establish. As a rule only a very severe blow to the chest wall produces a detectable cardiac injury. Many boxers and victims of accidents who sustain injuries to the chest show no anatomical or electrocardiographic evidence of myocardial damage.^{44, 100}

Many cases of alleged traumatic coronary occlusion reported in the literature probably represent instances of the coincidental occurrence of coronary occlusion and trauma. The electrocardiograms which had been reported in a large number of these patients as showing an occlusion actually showed only ST and T wave changes which occur in coronary insufficiency and not in occlusion.^{106, 107} In some instances a careful history will elicit the fact that the precordial pain preceded the accident. Although we are certain that episodes of coronary insufficiency may be precipitated by effort or trauma we believe that the occurrence of coronary occlusion is coincidental with extreme physical exertion and that the occlusion had occurred prior to the injury.^{17, 173} Trauma to the chest wall occasionally may

produce subendocardial tears and myocardial contusions in otherwise normal hearts ¹²⁰ All cases of so called steering wheel accidents should be carefully observed for several days in a search for electrocardiographic or clinical evidence of heart damage even in the absence of rib fracture or severe external injury ⁴⁷ These patients may show various arrhythmias without other changes for several days following the accident All cases with a definite history of severe chest trauma that develop precordial pain or discomfort at the time of the injury should be suspected of having sustained a myocardial contusion the patients should therefore be put to bed and be given adequate sedation

The following criteria help one in determining whether the cardiac damage was caused by a specific trauma ¹⁴⁷ The decision must be made on the basis of the facts in each case but in general we require the presence of two or more of these criteria for causal relationship

- 1 Significant electrocardiographic changes must be present
- 2 Cardiac enlargement unexplained by pre existent cardiac disease or hypertension must appear
- 3 Abnormal rhythms not including premature beats unless they are accompanied by other electrocardiographic changes must occur
- 4 A pericardial friction rub must be heard or evidence of effusion found
- 5 Congestive heart failure precipitated by the blow or strain must develop
- 6 Clinical signs and laboratory evidence of coronary insufficiency and or myocardial infarction must become manifest within twenty four hours following the incident

Such patients should be treated in the same manner as patients with nontraumatic coronary insufficiency or occlusion Anticoagulant drugs should be withheld however

because of the frequent occurrence of pericardial hemorrhage. If hemopericardium does occur the blood should be repeatedly aspirated if necessary.

Rupture of the heart occurs in a few of these cases within ten days to two weeks after the trauma but if the patient is carefully treated and put to bed as soon as symptoms appear, the incidence of this fatal complication may be reduced.

Many cases of traumatic myocardial damage recover completely after three to five weeks of bed rest, and show no clinical or electrocardiographic evidence of heart disease for years after the accident.

Chapter

9

SURGICAL EMERGENCIES IN CARDIOVASCULAR DISEASE THEIR PREVENTION AND TREATMENT

Surgery in the Cardiac Patient —The need for surgical treatment of an intercurrent disease among cardiac patients often arises. This need has increased in frequency with the increase in the number of the aged many of whom suffer from some cardio vascular disorder. Careful medical supervision of such patients is necessary in order to prevent the development of acute congestive failure coronary insufficiency tachycardias and pulmonary embolism following operation.

Most of the patients who have heart disease tolerate an operation well the exceptions being those who are in intractable failure or are recovering from a recent myocardial infarction. Serious complications do not develop even in these patients if the pre and postoperative treatment is properly administered.

A seventy eight year-old man who was studied because of increasing difficulty in urination exemplifies this point. A firm enlarged prostate was found and a prostatectomy was suggested but the patient was in severe congestive heart failure. He was rapidly digitalized and placed on rigid salt restriction. A complete A V block with a ventricular rate of 36 developed and the patient experienced several Stokes Adams seizures until the digitalis was stopped. The danger involved in the operation was ex-

plained to the patient, but he insisted on its being performed. During the operation profuse hemorrhage occurred and the patient went into mild shock. A continuous transfusion was administered together with repeated small doses of ephedrine and neosynephrine intramuscularly. The patient received 2500 cc. of blood within twenty four hours without developing congestive failure. There were no further Stokes Adams seizures. He made an uneventful recovery, and has remained well during the past four years.

Age should not be a deterring factor in patients who require surgery.²⁰⁴ Many elderly patients with heart disease and resultant severe cardiac disability, moreover are improved by the surgical removal of foci of infection and irritation. M. S., a fifty nine year-old housewife who had been an invalid because of severe recurrent attacks of angina pectoris, also suffered from biliary colic. The surgeons were hesitant about operating on her because of her cardiac condition but they were finally prevailed upon to do so. She tolerated the operation very well and was then practically free of her anginal attacks for two years. M. R., a sixty year old woman with gallbladder disease suffered an acute posterior wall infarction. Her convalescence was marked by repeated attacks of biliary colic. A cholecystectomy was performed one month after the occlusion had occurred. Her postoperative course was uneventful and since then she has not complained of pain.

In the presence of severe heart failure or after a recent coronary occlusion an operation should be performed only when it is urgently needed, and only after careful preparation, if time permits. Even when proper preparation is not possible the results have often been good. When congestive failure exists rapid intravenous digitalization should be carried out and mercurial diuretics should be given. In patients with rheumatic heart disease and auricular fibrillation the rate should be slowed by intravenous digitalis therapy before surgery is attempted. Persons with compensated heart disease may be expected to

undergo operations as well as those whose hearts are normal if certain precautions are taken. Special attention should be paid to the amount of fluids and sodium given pre and postoperatively and an adequate quantity of oxygen should be administered during the operation. The administration of large quantities of intravenous saline after an operative procedure should be avoided in cardiac patients. The measures suggested reduce the incidence of pulmonary edema and congestive heart failure.

In patients with coronary artery disease particular care should be exercised to avoid anoxemia especially during the induction of anesthesia. The most desirable anesthetic of course, is a local or regional one. Spinal anesthesia has usually been considered dangerous for patients with coronary artery disease or severe hypertension but we have found it effective and safe for operations on the lower abdomen or pelvis. Intramuscular neosynephrine or ephedrine may be given prophylactically to prevent a precipitous fall in blood pressure during spinal anesthesia and intravenous neosynephrine (0.5 to 1 mg) should be available for immediate use if this should occur.

If a general anesthetic is used ether is preferred or a combination of pentothal, oxygen and cyclopropane may be used. Since the latter may induce ventricular arrhythmias¹³¹ quinidine two doses of 6 gr (4 gm) orally. Pronestyl¹³⁴ (1 to 2 gm) or Dibenamine (7 to 7.5 mg/kg)¹³³ should be given before the operation particularly in patients with a history of arrhythmias. These agents markedly reduce the incidence of tachycardias that occur during surgical procedures. If auricular tachycardia flutter or fibrillation occurs quinidine should be given intramuscularly. If a ventricular arrhythmia sets in despite these prophylactic measures Pronestyl should be injected intravenously.

The incidence of postoperative thromboembolism and pulmonary infarction is higher in cardiac than in non cardiac patients. The single most important factor in preventing these complications is *early ambulation, par*

ticularly in the obese. The patient should also be instructed to turn in bed as soon as possible to move his legs within several hours after operation, and to avoid straining at stool. If pulmonary infarction occurs, despite these measures, treatment is given as outlined in Chapter 3 p. 83.

Cardiac arrest may occur during the course of any operative procedure in both normal and cardiac patients. It may be due to marked vagal stimulation during the operation and can be prevented in some cases by the intravenous administration of atropine, 0.6 mg (1/100 gr), sixty minutes before surgery, or 0.3 mg (1/50 gr) intramuscularly thirty minutes before.¹²² Cardiac arrest occurs in operations on the chest more often than in abdominal or pelvic surgery. When it does occur, the instantaneous execution of a well planned course of treatment is essential to save the patient's life.

A sixteen year old boy, who was operated upon for atelectasis of the left lung secondary to fibrosarcoma of the bronchus, furnishes a dramatic example of the success of cardiac resuscitation, even after a forty minute period of cardiac arrest.* He received cyclopropane and ether at 8:10 A.M., at 8:25 A.M., an oval endotracheal tube was introduced. At 8:27, the left hemithorax was entered. Sixty mg of procaine HCL was injected into the tubing of an intravenous drip. The pulmonary artery was ligated at 9:05, when cardiac arrest occurred. Artificial respiration was begun immediately by intermittent manual compression of the breathing bag of the anesthetic apparatus. Simultaneously, manual massage of the heart was started at a rate of 50 per minute. At 9:10, a mixture of 80 mg of procaine HCL and 1 mg epinephrine was injected into the left ventricular cavity. At 9:25, 60 mg of procaine and 0.5 mg epinephrine were given and at 9:30, 80 mg more of procaine were injected. Cardiac action did not return. At 9:45, the pericardium was grasped with a hemostat,

* This case is described with the permission of the operating surgeon Dr. Arthur S. W. Touroff.¹²³

preparatory to incision. At that moment the heart suddenly beat violently and normal pulsation and respiration began. The pneumonectomy was completed without mishap. On the twelfth postoperative day the patient was entirely normal and has remained well.

Restoration of heart action and maintenance of an adequate oxygen supply to the tissues are the two main objectives of therapy in these cases. The first step is to expose the heart as quickly as possible preferably directly through the chest. Sterile precautions are not required and the procedure may be carried out by anyone available. Upon exposure of the heart cardiac arrest is usually seen but in some cases the collapse is the result of ventricular fibrillation. If the latter is found 5 cc. of a 1 per cent Procaine solution should be injected into the right auricle and the epicardial surface of the heart should be bathed by 2 to 3 cc.¹⁴⁸ in preparation for defibrillation. This is accomplished by placing electrodes on either side of the heart and passing a current of 110 volts and 1.5 amperes through it.¹⁴⁹ When cardiac standstill has thus been produced the treatment already described should be instituted. Manual cardiac massage is started and continued at a steady rate of 50 to 80 per minute¹⁵⁰ until spontaneous cardiac action begins. If there is no response 0.5 to 1 cc. of a 1:1000 solution of epinephrine is injected into the right auricle.¹⁵¹ If the arrest still persists the massage and epinephrine injections should be continued for at least forty to sixty minutes before admitting failure. When the heart is unusually flabby or atonic 3 to 5 cc. of a 1 per cent solution of calcium chloride may be effective.

Artificial respiration should be started simultaneously with the exposure of the heart if respirations have ceased. If possible an intratracheal tube should be inserted and the lungs inflated and deflated by mechanical means. This is by far the most efficient method of supplying oxygen.¹⁵² If endotracheal tubes are not available the resumption of respiration may be effected by mouth to mouth breathing.

alternating manual pressure and release of the costal cage or the use of a tight fitting face mask pressure being applied intermittently. When the latter method is used there is a real danger of forcing air into the stomach and causing marked gastric dilatation.²⁷³ One hundred per cent oxygen should be given and CO₂ mixtures should be avoided since the patient already has accumulated an excessive amount of CO₂ in the tissues. Adequate oxygenation of the brain and other organs may be maintained by use of any one of the above procedures until spontaneous respiration and circulation are resumed.

When hemorrhage has occurred and the blood volume is markedly decreased intra arterial transfusions of blood should be given at the rate of 150 to 200 cc minute. These transfusions are given at a pressure higher than the systolic blood pressure.

In many cases patients who had apparently expired will be completely revived, and will recover with no ill effects if the suggested treatment is undertaken promptly. If the surgeon is properly equipped and psychologically ready for this emergency, good results may be expected.

BIBLIOGRAPHY

- 1 ABBOTT K H Acetylcholine in Paroxysmal Tachycardia
J A M A 113 1243 1939
- 2 ALTANA G Singular Electrocardiographic Picture Caused by
Circulation 1 28 1950
- 3 ANDI Fibrillation and
8 128 1932
- 4 ANREP G V DARSCH G S LILLY J and MISRAH J
J Amm Visnaga in the Treatment of the Anginal Syndrome
Brit Heart J 8 171 1946
- 5 ARMERLST C A and LEVINE S A Paroxysmal Ventricular
Tachycardia A Study of One Hundred and Seven Cases
Circulation 1 28 1950
- 6 ARONSTEIN C G and NELMAN L Ball Thrombus of Heart
Report of Case With Review of Literature Arch Path 27
907 1939
- 7 ASTWOOD E B and VANDERLAAN W P Treatment of Hyper
thyroidism With Propylthiouracil Ann Int Med 25 813
1946

- 8 ATKINSON W J JR Tetraethyl Ammonium Chloride in the Treatment of Angina Pectoris *Am Heart J* 39 336 1950
- 9 BARACH A L and MARTIN J Effect of Inhalation of Oxygen Under Positive Pressure on Pulmonary Congestion and Edema *J Clin Invest* 15 466 1936
- 10 BARBER H Trauma of Heart *Brit M J* 1 433 1938
- 11 BARKER N W HINES E A KVALF W F and ALLEN E V Dicoumarol Its Action Clinical Use and Effectiveness as an Anticoagulant Drug *Am J Med* 3 637 1947
- 12 BARONOFSKY D and SKINNER A Ligation of Left Auricular Appendage for Recurrent Embolism *Surg* 27 848 1950
- 13 BARROW J G Treatment of Paroxysmal Supraventricular Tachycardia with Lanatoside C *Ann Int Med* 32 116 1950
- 14 BATTERMAN R C and DEGRAFF A C Comparative Study on Use of Purified Digitalis Glycosides Digoxin Digitoxin and Lanatoside C for the Management of Ambulatory Patient with Congestive Heart Failure *Am Heart J* 31 663 1947
- 15 BATTERMAN R C UNTERMAN D and DEGRAFF A C The Subcutaneous Administration of Mercaptomerin (Thiomerin) Effective Mercurial Diuretic for Treatment of Congestive Heart Failure *J A M A* 140 1268 1949
- 16 BATTERMAN R C DEGRAFF A C and SHORR H M Further Observations on the Use of Mercupurin Administered Orally *Am Heart J* 31 431 1946
- 17 BECK C S Further Observations on Stab Wounds of Heart *Ann Surg* 115 698 1942
- 18 BECK C S STANTON E BLINCHOK W and JETER C Revascularization of Heart by Graft of Systemic Artery into Coronary Sinus *J A M A* 137 436 1948
- 19 BECK C S Resuscitation of Cardiac Standstill and Ventricular Fibrillation Occurring During Operation *Am J Surg* 54 273 1941
- 20 BECK C S PRITCHARD W H and FEIL H Ventricular Fibrillation of Long Duration Abolished by Electric Shock *J A M A* 135 985 1947
- 21 BELL G O BRADLEY R B and HURNTHAL L M Paroxysmal Tachycardia Experiences with Massive Doses of Quinidine Intravenously in a Refractory Case *Circulation* 1 919 1950
- 22 BELL G I BELL R E and WILSON D R Rheumatic Pancarditis Treated with Cortisone *Canad Med Assn J* 63 63 1950
- 23 BELLET S and NADLER C S Use of Quinidine Sulfate in the Treatment of Hiccup *Am J of Med Sc* 216 680 1948
- 24 BELL R A BELL R S and MASSIE F An Antidote

- 26 BIGGER I A Heart Wound Report 111
Upon in Medical College of Virginia Hospital
of Treatment and Prognosis J Thorac Surg
- 27 BLAND P F and SWEET R H A Venous Shunt
Mitral Stenosis JAMA 140 1259 1941
- 28 BLOOMFIELD A L Treatment of Atrial Fibrillation
Editorial Am J Med 437 1949
- 29 BLUMGART H L The Reaction to Exercise in the Heart
by Atrial Fibrillation Heart 11 49 1934
- 30 BLUMGART H L FEYEL S A and BERNIS D D
Heart Failure and Angina Pectoris The Therapeutic
of Thyroidectomy on Patient With Atrial Fibrillation
Evidence of Thyroid Toxicity Arch Int Med 57 506 1936
- 31 BLUMGART H L FREEDBERG A S and KILIAN C S Heart
thyroidism Produced by Radioactive Iodine in the
ment of Euthyroid Patients with Angina Pectoris
gestive Heart Failure (circulation 1 1105 1950)
- 32 BORG J F Treatment of Edema with Mercurial
Mercurial Diuretic Am Heart J 24 41 1947
- 33 BOYD L J and SCHERF D Magnesium Sulfate in Paroxysmal
Tachycardia Am J Med Sci 206 43 1943
- 34 BRADLEY S and EGLESTON C Postural Hypertension Am
Heart J 1 73 1925
- 35 BRANNON E S Hemiplegia Following Coronary Artery
Occlusion Case Report Am Heart J 36 299 1948
- 36 BRANNON E S STEAD F A JR WARRIN J A and MERRILL
A J Hemodynamics of Acute Hemorrhage in Man
Heart J 31 407 1946
- 37 BRILL I C Congestive Heart Failure Arising from Uncontrolled
Atrial Fibrillation in the Otherwise Normal Heart Follow-up
Notes on Previously Reported Case Am J Med 2 44
1947
- 38 BRILL I C and MEISSNER W A Role of Coronary Artery
Disease in Etiology of Atrial Fibrillation Ann Int Med
14 1341 1941
- 39 BURKE G F and WRIGHT I S Tromexan 33 Carboxy
methylenebis (4 Hydroxycoumarin Ethyl Ester) Experimental
and Clinical Properties Circ 3 164 1951
- 40 BERNSTEIN C Treatment of Acute Arrhythmias During Anes-
thesia by Intravenous Procaine Anesthesiology 7 113 1946
- 41 BURWELL W B and HENDRIX J P Digitalis Poisoning Am
J Med 8 640 1950
- 42 BURWELL C S and DIXTER L Benign Heart Disease for
A Amer Physicians 60 59 1947
- 43 BUTTERWORTH J S and POWELLER C A The Esophageal
Electrocardiogram in Arrhythmias and Tachycardia Am
Heart J 32 681 1946
- 44 ----- Electrocardiographic Study of the Effects of Boxing
Am Heart J 23 49 1942

- 45 COHEN S M EDHOLM E G HOWARTH S McMICHAEL J
and SHARPEY SCHAFFER E P Cardiac Output and Periph-
eral Blood Flow in Arteriovenous Aneurysm *Clin Sc* 7 35
1948
- 46 COHN A E and STEWARD H J Evidence That Digitalis
Influences Contraction of the Heart in Man *J Clin Invest*
1 97 1924
- 47 COMFAU W J The Mechanism of Syncopal Attacks Associated
with Paroxysmal Auricular Fibrillation *New England J Med*
227 134 1942
- 48 CONN J W The Diagnosis and Management of Spontaneous
Hypoglycemia *JAMA* 134 140 1947
- 49 COOK L Recovery After Massage of the Heart *British Med J*
2 1118 1926
- 50 COOKSON H Aetiology and Prognosis of Auricular Fibrillation
Quart J Med 23 309 1930
- 51 Cornell Conferences on Therapy Management of Disorders of
Cardiac Rhythm *Am J Med* 5 110 1948
- 52 Cornell Conferences on Therapy Use of Mercurial Diuretics
New York State J Med 46 62 1946
- 53 DACK S MASTER A M HORN H GRISHAM A and FIELD L
Acute Coronary Insufficiency Due to Pulmonary Embolism
Am J Med ~ 464 1949
- 54 DAVIES D T MANSELL H E and O'SHAUGHNESSY L Sur-
gical Treatment of Angina Pectoris and Allied Condition
Lancet 1 1 1938
- 55 DEARING W H BARNES A E and ESSEX H E Myocardial
Lesions Produced by Digitalis in the Presence of Hyperthy-
roidism *Circulation* 1 394 1950
- 56 DECHARD G M JR and HERMANN G R Association of
Paroxysmal Atrial Tachycardia with Atrial Flutter or Fibrilla-
tion *Am Heart J* 28 457 1944
- 57 DEGRAFF A C Clinical Comparison of Cardiac Glycosides
New York State J Med 46 1803 1946
- 58 DEGRAFF A C and BATTERMAN R C Recent Advances in
Digitalis Therapy with Particular Attention to the Use of
Pure Glycosides *M Clin North America* 26 929 1942
- 59 DEROW H A and WOLFF I Oral Administration of Meru-
purin Tablets in Ambulatory Patients with Chronic Congestive
Heart Failure *Am J Med* 3 693 1947
- 60 DOCK W and TAINTER M L The Circulatory Changes After
Full Therapeutic Doses of Digitalis with a Critical Discussion
of Views on Cardiac Output *J Clin Investigation* 8 467
1930
- 61 DOCK W Sodium Depletion as a Therapeutic Procedure The
Value of Ion Exchange Resins in Withdrawing Sodium from
the Body *Tr A Am Physicians* 59 282 1946
- 62 DOCK W (quoted in reference 165)

- 63 EDHOLM O G HOWARTH S and McMICHAEL J Heart Failure and Bone Blood Flow in Osteitis Deformans. Clin Sc 5 249 1945
- 64 EDHOLM O G Effect of Gravity on the Blood Pressure of the Cat J Physiol 98 79 1940
- 65 EDELL G A and ...
- 66
- 641 1944
- 67 EICHNA L W TALBE H and DeGRAFF A C Serial Deter...
- 68 ELLIS I B and HAYNES F W Postural Hypotension With Particular Reference to its Occurrence in Disease of the Central Nervous System Arch Int Med 58 773 1936
- 69 ELLIS L B Studies in Complete Heart Block Am J Med Sc 183 225 1932
- 70 ENGEL G L Mechanism of Fainting J of Mt Sinai Hosp 12 170 1945
- 71 ENGLEMAN E P KRUPP M A and KUNKEL P The Oral Use of Cortisone Suspension in Syrup J A M A 145 402 1951
- 72 EISENBERG C D SIMMONS H G and MINTZ A A Effects of Potassium Upon the Heart with Special Reference to Possibility of Treatment of Toxic Arrhythmias Due to Digitalis Am Heart J 39 713 1950
- 73 EPSTEIN F H and RELMAN A Transfusion Treatment of Shock Due to Myocardial Infarction New England J Med 241 889 1949
- 74 ERNSTEN A C Cardiovascular Complications of Hyperthyroidism Am J Med Sc 195 248 1938
- 75 EVANS W Unity of Paroxysmal Tachycardia and Auricular Flutter Brit Heart J 6 221 1944
- 76 EVANS W DICK P and EVANS B Rapid Digitalization Brit Heart J 10 103 1948
- 77 EVANS W and BENSON R Mass Thrombus of Left Auricle Brit Heart J 10 39 1948
- 78 FAUTEX M J Cardiac Resuscitation Thoracic Surg 16 623 1947
- 79 FEIL H S and GILDER M D The Regularity of Simple Paroxysmal Tachycardia Lond Heart 8 1 1921
- 80 FERRER M I HARVEY R M WELSH L DRESDALE D T COURVAND A and RICHARDS D W Some Effects of Quinidine Sulfate on the Heart and Circulation in Man Am Heart J 36 816 1948

- 81 FERRER M I and SOKOLOFF L Antidiuretic Effect of Morphine and Demerol in Congestive Heart Failure *Am J of Med Sc* 214 372 1947
- 82 FOLEY W T and WRIGHT I S Long Term Anticoagulant Therapy for Cardiovascular Disease *Am J Med Sc* 217 136 1949
- 83 FOX S L Radioiodine in the Management of Thyrocardiac Disease *Ann Western Med and Surg* 4 282 1950
- 84 FRIEDBERG C K Diseases of the Heart W B Saunders and Co Phila 1950 p 240
- 85 ——— (ibid) p 274
- 86 ——— (ibid) p 494
- 87 FRIEDBERG C K and HORN H Acute Myocardial Infarction Not Due to Coronary Artery Occlusion *J A M A* 112 1675 1939
- 88 FRIEDBERG C K Revised Criteria for Diagnosis and Treatment of Subacute Bacterial Endocarditis *Med Clin N Y* 769 (May) 1950
- 89 FRIEDLANDER R D and IRVING S A Auricular Fibrillation and Flutter without Evidence of Organic Heart Disease *New England J Med* 211 624 1934
- 90 FRIEDMAN M and BINI R JR Observation Concerning the Influence of Potassium Upon the Action of a Digitalis Glycoside (Janatoside C) *Am J Med Sc* 214 633 1947
- 91 GAILLARDIN L Tachycardia Paroxystique Ventriculaire avec conservation du Rythme Auriculaire normal caractères du pouls Reineux Enregistrement de la Terminaison de l'acces *Arch d mal du coeur* 13 121 1920
- 92 GARVIN C I Infarction in Heart Disease *Am J M Sc* 203 473 1942
- 93 GERTLER M M and YOHALIM S B Effect of Atrabrine on Auricular Fibrillation and Supraventricular Tachycardia in Man *J Mt Sinai Hosp* 13 323 1947
- 94 GLUCK J GOLD H GREIFER T MODFELL W KWIT N T THICKMAN S OTTO H I and WARSHAW L Quinidine Sulfate in Propylene Glycol by Intramuscular Injection in Man *J A M A* 145 637 1951
- 95 GOLD H Quinidine in Disorders of the Heart Paul B Hoeber Inc N Y 1950
- 96 ——— (ibid) p 13
- 97 GOLD H and CATTELL M Mechanism of Digitalis Action in Abolishing Heart Failure *Arch Int Med* 65 263 1940
- 98 GOLDENBERG M SNYDER C H and ARANOW H JR New Test for Hypertension Due to Circulating Epinephrine *J A M A* 135 971 1947
- 99 GONZALEZ SARATHI L On the Intravenous Use of Morphine in the Treatment of Paroxysmal Ventricular Tachycardia *Am Heart J* 33 719 1947

- 100 GORE IRA The Question of Traumatic Heart Disease Ann Int Med 33 865 1950
- 101 GORHAM L W LESTER D E WOLF A V and SHULTZ H H Relative Importance of Dietary Sodium Chloride and Water Intake in Cardiac Edema Ann Int Med 27 575 1947
- 102 GORLIN R SAWYER C G HAYNES F W GOODALE W T and DEXTER L Effects of Exercise on Circulatory Dynamics in Mitral Stenosis III Am Heart J 41 192 1951
- 103 GOSSAGE A M and HICHS J A B On Auricular Fibrillation Quart J Med 6 435 1912
- 104 GREINER T GOLD H CATTELL M TRAVELL J BAASST H RINZLER S H BENJAMIN Z H WARSHAW L F BOBB A L KIVIT N T MODELL W ROTHENDLER H H MESSELOFF C R and KRAMER M L A Method for the Evaluation of Effects of Drugs on Cardiac Pain in Patients with Angina of Effort A Study of Khe'll'n (Vitamin) Am J Med 9 143 1950
- 105 GROSSMAN J WESTON R E EDELMAN I S and LEITER I Studies on Thimerin—A Subcutaneously Administerable Mercurial Diuretic Circulation 1 508 1950
- 106
- 107
tensive and Hypertensive Subjects. Proc Soc Exper Biol and Med 27 477 1951
- 108 HARKEN D E and WILLIAMS A C Foreign Bodies in and in Relation to Thoracic Blood Vessels and Heart Migratory Foreign Bodies Within Blood Vascular System Am J Surg 72 80 1946
- 109 HARKEN D E Review of Activities of Thoracic Center for III and IV Hospital Groups 160th General Hospital European Theatre of Operations June 10 1944 to Jan 1 1945 J Thoracic Surg 15 31 1946
- 110 HARKEN D E ELLIS L B WARE P F and NORMAN L R The Surgical Treatment of Mitral Stenosis Valvuloplasty Circulation 22 226 231 1951
- 111 HAL
WINSON O T
Evaluation of
Study of the
Radioactively Labeled Anticoagulants in Animals Circ J 171 1951
- 112 HELLERSTEIN H K and MARTIN J W Incidence of Thrombo-Emboic Lesions Accompanying Myocardial Infarction Am Heart J 33 443 1947
- 113 HENCH P S SLOCUM C H BARNES A R SMITH H L

- 114 HENDERSON V and PRINCE A The Oxygen Pulse and the Systolic Discharge *Am J of Phys* 25 106 1914
- 115 HERRMANN G R and HEJTMANCIK M R A Clinical and Electrocardiographic Study of Paroxysmal Ventricular Tachycardia and Its Management *Ann Int Med* 28 989 1948
- 116 ——— Atrial Flutter II Methods of Treatment *Am Heart J* 41 182 1951
- 117 HESSE E Ueber die Funktionsfähigkeit des Verletzten und genaheten Herzens im Lichte der Dauerresultate der Herznaht *Deutsche Ztschr f Chir* 190 239 1925
- 118 HILDEN T On Pathogenesis of Acute Pulmonary Edema *Acta Med Scand Supp* 234 162 1949
- 119 HOCK G A Pheochromocytoma with Paroxysmal Hypertension *J Urology* 61 473 1949
- 120 HODGES H A and GILMOUR M T Myocardial Contusion Report of a Fatal Case with Autopsy *Ann Int Med* 34 248 1951
- 121 HOMANS J Venous Thrombosis in Lower Limbs Its Relation to Pulmonary Embolism *Am J Surg* 38 316 1937
- 122 HORN H and FINKELSTEIN J E Arteriosclerosis of Coronary Arteries and Mechanism of their Occlusion *Am Heart J* 19 655 1940
- 123 HOWARTH S McMICHAEL J and SHARPEY SCHAFFER E P Effects of Venesection in Low Output Heart Failure *Clin Sc* 6 41 1946
- 124 HOY S H and WOOD J F JR Cation Exchange Resins in the Treatment of Congestive Heart Failure *Ann Int Med* 33 1139 1950
- 125 IDF I N The Clinical Aspects of Complete Auriculoventricular Heart Block Clinical Analysis of 71 Cases *Ann Int Med* 32 510 1950
- 126 Inhalation Therapy *JAMA* 144 25 1950
- 127 JAFFE H MASTER A M and DORRANCE W R The Salt Depletion Syndrome Following Mercurial Diuresis in Elderly Persons *Am J Med Sc* 220 60 1950
- 128 JAFFE H L Electrocardiographic Changes of Coronary Insufficiency During the Formative Stage of Coronary Occlusion *J Mt Sinai Hosp* 16 404 1950
- 129 JAMES D F TURNER H and MERRILL J Circulatory and Renal Effects of Aminophyllin in Congestive Heart Failure *Am J Med* 5 619 1948 Ab
- 130 JOHNSON J and KIRBY C K Cardiac Resuscitation *S Clin North America* 29 1745 1949
- 131 JOHNSTON M Cyclopropane Anesthesia and Ventricular Arrhythmias *Brit Heart J* 12 239 1950
- 132 ——— General Anesthesia and Cardiac Inhibition *Brit Heart J* 13 47 1951

- 133 JORDAN R A SCHEFFLEY C H and EDWARDS F F Mural Thrombosis and Arterial Embolism in Mitral Stenosis A Clinico-Pathologic Study of 51 Cases *Circulation* 3 363 1951
- 134 JOSEPH S I ROVENSTINE E A Cardiac Arrhythmias Prophylaxis and Therapy with Procaine Amide During Cyclopropane Anaesthesia for Thoracic Surgery *J Pharm and Exper Ther* 109 19 1951 Ab
- 135 KATZIN H M WALLER J V and BLUMGART H L Cardiac Cirrhosis of the Liver *Arch Int Med* 64 457 1939
- 136 KAUFMAN R E Immediate Fatalities after Intravenous Mercurial Diuretics *Ann Int Med* 28 1040 1948
- 137 KAYDON H J STEELE J M MARK L C and BRODIE B B The Use of Procaine Amide in Cardiac Arrhythmias *Circulation* In press
- 138 KELLEY J L Saddle Embolism of the Aorta Report of Successful Embolectomy *Ann Surg* 128 257 1948
- 139 KEITH V M BARRIER C W and WHELAN M The Diuretic Action of Ammonium Chloride and Novasurol *J A M A* 85 799 1925
- 140 KELLY H G and BAYLESS R J S Influence of Heart Rate on Cardiac Output *Lancet* 2 1071 1949
- 141 KISCH B Das Vorkommen und der Nachweis von Interpolierten Vorhofextrasystolen (Zugleich ein Beitrag zur Kenntnis der Partial extrasystolen) *Ztschr f d ges Exp Med* 25 188 1921
- 142 ——— Strophanthin Brooklyn Medical Press 1944
- 143 KISSANE R W BROOKS R and CLARK T Relation of Supraventricular Paroxysmal Tachycardia to Heart Disease and the Basal Metabolism Rate *Circulation* 1 950 1950
- 144 KISSANE R W and KOONS R A Spontaneous Redigitalization Following Rapid Diuresis in Congestive Heart Failure *Tr Am Therap Soc* 39 111 1940
- 145 KOSSMAN C E and BERGER A R Auricular Flutter of 11 Years Duration with Observations on Esophageal Electrocardiograms *Ann Int Med* 15 128 1941
- 146 KOLATZ N B SMITH J R and WRIGHT S T Observations on the Effect of Tourniquets on Acute Cardiac Crises Normal Subjects and Chronic Heart Failure *Am Heart J* 23 624 1942
- 147 KULKA W Contusion of the Heart *Am Heart J* 39 438 1949
- 148 LADLE J S MURISON P J and PACK C T Use of Tetraethyl Ammonium Bromide as a Diagnostic Test for Pheochromocytoma *Proc Am Federation Clin Research* 3 67 1947
- 149 LEVY C M STRAZZA J A and JAFFIN A E Fluids in Con

- 151 LEVINE S A Some Harmful Effects of Recumbency in Treatment of Heart Disease *J A M A* 126 80 1944
- 152 LEVINE S Clinical Heart Disease W B Saunders Co 1945 pg 201
- 153 LEVINE S A Oculocardiac Reflex Electrocardiographic Study With Special Reference to Differences Between Right and Left Vagal and Ocular Pressures in Tibetics and Non tibetics *Arch Int Med* 15 758 1915
- 154 LEVINE S A and STEVENS W B The Therapeutic Value of Quinidine in Coronary Thrombosis Complicated by Ventricular Tachycardia *Am Heart J* 3 253 1928
- 155 LEVINSON D C EDMANDES D T and GRIFFITH G Dissecting Aneurysm of the Aorta Its Clinical Electrocardiographic and Laboratory Features *Circulation* 1 360 1950
- 156 LEWIS T The Mechanism and Graphic Registration of the Heart Beat Shaw and Sons Ltd London 4th Ed 1925
- 157 ——— Observations Upon Disorders of the Heart's Action *Heart* 3 279 1911 12
- 158 ——— Lecture on Vasovagal Syncope and Carotid Sinus Mechanism with Comments on Gowers and Nothnagel's Syndrome *Brit M J* 1 873 1932
- 159 LINFenthal A J and FREEDBERG A S Medical Progress Measures Used in the Prevention and Treatment of Cardiac Arrhythmias *New Eng J Med* 241 570 1949
- 160 LONGMIRE W P BEAL J M and LEAKE W N Resection of the Auricular Appendages *Diseases of the Chest* 19 307 1951
- 161 LUISADA A Pathogenesis of Paroxysmal Pulmonary Edema *Medicine* 19 475 1940
- 162 LUNDY C J and McLELLAN L I Paroxysmal Ventricular Tachycardia An Etiological Study with Special Reference to the Type *Ann Int Med* 7 812 1934
- 163 MACGUIRE W B JR Risk of Uremia Due to Sodium Depletion *J A M A* 137 1377 1948
- 164 MACLEAN A R ALLEN E V and MAGATH T B Orthostatic Tachycardia and Orthostatic Hypotension Defects in Return of Venous Blood to the Heart *Am Heart J* 27 145 1944
- 165 MADDEN J I Resection of Left Auricular Appendix A Prophylaxis for Recurrent Arterial Emboli *J A M A* 140 769 1949
- 166 MANCHESTER B and RABKIN B Acute Coronary Occlusion Treated With Dicoumarol and Regulated by a Simple Bed side Blood Prothrombin Determination In Preparation
- 167 MARK L C BERLIN I KAYDON H J ROVENSTINE E A STEFFLE J M and BRODIE B B The Action of Procaine Amide on Ventricular Arrhythmias *J Pharm and Exper Ther* 98 21 1950

- 168 MARSHALL D S Recognition and Emergency Treatment of Venous Thrombosis and Its Complications Med Clin of N A Nov 1949
- 169 MARVIN H M Paroxysmal Ventricular Tachycardia with Alternating Complexes Due to Digitalis Intoxication Am Heart J 4 21 1928
- 170 MASSELL B F and WARREN J E Effect of Pituitary Adrenocorticotrophic Hormone (ACTH) on Rheumatic Fever and Rheumatic Carditis JAMA 144 1335 1950
- 171 MASSELL B F WARREN J E STURGIS G P HALL B and CRAIGE E The Clinical Response of Rheumatic Fever and Acute Carditis to ACTH New England J of Med 242 692 1950
- 172 MAS - - - - -
- 173 MAS - - - - -
359 1940
- 174 MASTER A M DACK S and JAFFE H L Disturbances in Rate and Rhythm in Acute Coronary Artery Thrombosis Ann Int Med 11 735 1937
- 175 - - - - - Partial and Complete Heart Block in Acute Coronary Artery Occlusion Am J Med Sci 190 513 1938
- 176 MASTER A M GUBNER R DACK S and JAFFE H L Differentiation of Acute Coronary Insufficiency with Myocardial Infarction from Coronary Occlusion Arch Int Med 67 647 1941
- 177 MASTER A M JAFFE H L DACK S and GRISHMAN A Coronary Occlusion Coronary Insufficiency and Angina Pectoris Am Heart J 27 803 1944
- 178 MASTER A - - - - - and HORN H Acute Hemorrhagic Product
- 179 MASTER A - - - - -
ular Fit
- 180 MASTER A M Digoxin Intoxication J
- 181 - - - - - Treatment of Cardiovascular Emergencies U S Nav M Bull 39 100 1941
- 182 MASTER A M JAFFE H L and DORRANCE W R The Treatment of Heart Failure New York State J of Med 50 553 1950
- 183 - - - - -
- 184 - - - - -
caine Amide Hydrochloride in Supraventricular Arrhythmias Circ 4 387, 1951

- 185 MANS J H Dyspnoea *Medicine* 3 309 1924
- 186 MERRILL A J Edema and Decreased Renal Blood Flow in Patients with Chronic Congestive Heart Failure Evidence of Forward Failure as the Primary Cause of Edema *J Clin Investigation* 25 389 1946
- 187 MERRILL J P LEVINE H D SOMERVILLE W and SMITH S Clinical Recognition and Treatment of Acute Potassium Intoxication *Ann Int Med* 33 797 1950
- 188 MILLER G G and RIPSTEIN C B Resuscitation of the Surgical Patient *Canad M A J* 61 255 1949
- 189 MILLER R HARVEY W and FINCH C Antagonism of Dicoumarol by Vitamin K Preparation *New England J Med* 242 211 1950
- 190 MOKOTOFF R ROSS G and LEITER L Renal Plasma Flow and Sodium Reabsorption and Excretion in Congestive Heart Failure *J Clin Investigation* 27 1 1948
- 191 MOSER M Personal Observation
- 192 MOSER M SHEPHERD G and SCHWINGER H Pheochromocytoma with Calcification Simulating Cholelithiasis *Radiology* 55 855 1950
- 193 MURPHY F D COVILL H and GRILL J C The Effects of Intravenous Solutions on Patients With and Without Cardiovascular Defects *JAMA* 116 104 1941
- 194 NATHAN L H and HOFF H C Auricular Fibrillation in Hyperthyroid Patients Produced by Acetyl B Methyl Choline Chloride with Observations on Role of Vagus and Some Exciting Agents in Genesis of Auricular Fibrillation *JAMA* 105 254 1935
- 195 NATHANSON M H Further Observations on Effects of Drug on Induced Cardiac Standstill Effect of Epinephrine and Related Compounds *Arch Int Med* 54 111 1934
- 196 ———— Modification of Vagus Inhibition of Heart by Quinidine *Proc Soc Exper Biol and Med* 31 1234 1934
- 197 NATHANSON M H ENGELBERG H and HERSH J Cardiovascular Effects of Paredrine *Am Heart J* 24 153 1942
- 198 NICKERSON M BROWN H O and SCOTT M Mechanism and Efficacy of Dibenamine Protection Against Spontaneous Cyclopropane Arrhythmias in Surgical Patients *Am J Med* 6 396 1949
- 199 NICKERSON M and BROWN H O Dibenamine Protection Against Spontaneous Cyclopropane Arrhythmias in Surgical Patients *Anesthesiology* In Press
- 200 NIDELMAN P L LEFF I L and HOWE C D Thrombopenic Purpura Following Quinidine *JAMA* 137 1219 1948
- 201 OHLER H I KATZ K H and WAGNER D J Khellin in the Treatment of Angina Pectoris *New England J Med* 244 315 1951
- 202 PAINE R and SMITH J R The Mechanism of Heart Failure *Am J Med* 6 84 1949

- 203 PARKINSON J PAPP C and FRANK W The ECG in the Pathogenesis of Stokes-Adams Attack Brit Heart J 3 171 1941
- 204 PARONIS W H and PARKS W K The Elderly Patient and Surgical Risk Arch Surg 58 588 1949
- 205 PARNISH A E The Bioassay of Adrenal Cortex in the Treatment of Patients with Congestive Heart Failure J Clin Invest 28 45 1949
- 206 PATTERSON J C Relation of Physical Exertion and Precipitation of Coronary Thrombi JAMA 112 895 1949
- 207 PEARSON R S B Sinus Bradycardia with Cardiac Arrest Brit Heart J 7 85 1945
- 208 PHILLIPS E and LEVINE S A Atrial Fibrillation With and Without Other Evidence of Disease Am J Med 7 478 1949
- 209 PLOYZ M Acute Urinary Retention Following the Use of Mercaptopurine in Heart Disease Urol and Cut n Rev 47 786 1941
- 210 PRANDONI A and WRIGHT I S Anticoagulants Heparin and the Dicoumarin 3,3-Methylene-bis (4-hydroxy coumarin) Bull New York Acad Med 18 443 1942
- 211 PRINZMETAL M CORDAY E OBLATH R W BRILL I C and KRUEGER H L The Nature of Atrial Flutter in Man Proc Am Heart Assn June 1950
- 212 PRINZMETAL M CORDAY E BRILL I C SELLERS A I OBLATH R W FLEIG W A and KRUEGER H L Mechanism of the Atrial Arrhythmias Circulation 1 241 1950
- 213 RAAB W Propylthiouracil for Angina Pectoris Acta Med Scandinav 135 364 1949
- 214 RAVITCH M M and BLALOCK A Aspiration of Blood from Pericardium in Treatment of Acute Cardiac Tamponade After Injury Further Experience with Report of Cases Arch of Surg 58 463 1949
- 215 REVENO W S Thrombolysis in Angina Pectoris Am J Med 1 607 1946
- 216 RICE R L ACKERMAN J S and SAUBER R Long Term Dicoumarol Therapy Ann Int Med 32 735 1950
- 217 RICE T B Low Sodium Diet Philadelphia Lea & Febiger 1951
- 218 RICHARDS D W Jr Dynamic of Congestive Heart Failure Am J Med 6 772 1949
- 219 ——— Cardiac Output by the Catheterization Technique in Various Clinical Conditions Federation Proc 4 215 1945
- 220 RINZLER S H BANSY H BENJAMIN Z H BOBB A L and TRAVELL J Failure of Alpha-Tocopherol to Influence Chest Pain in Patients with Heart Disease Circulation 1 288 1950
- 221 ROBY W H Visual Disturbances with Digitalis Medication New England J Med 213 248 1945
- 222 ROTH G M and KVALE W F Tentative Test for Thrombocytopenia Am J Med Sc 210 631 1945
- 223 SIMPSON J J and SINGER I M Plasma and Blood Infusions Following Myocardial Infarction Am Heart J 36 54 1947

- 224 SCHERF D ROMANO F J and TERRANOVA R Experimental Studies on Auricular Flutter and Auricular Fibrillation *Am Heart J* 36 241 1948
- 225 SCHNITKER M A and LEVINE S A Presence of Digitalis in Body Fluids of Digitalized Patients *Arch Int Med* 60 240 1937
- 226 SCHWARTZ S P and BILOON S Clinical Signs of Occluding Thrombi of Left Auricle *Am Heart J* 7 84 1931
- 227 SCHWARTZ S P and DE SOLA POOL N Transient Ventricular Fibrillation *Am Heart J* 39 361 1950
- 228 SCHWARTZ S P and LEVY J Digitalis Studies on Children with Heart Disease Effects of Digitalis on Edema of Children With Rheumatic Fever and Chronic Valvular Heart Disease in Presence of Sinus Rhythm *Am J Dis Child* 42 1349 1931
- 229 SCHWARTZ W B and LEVINE S A Case of Paroxysmal Auricular Tachycardia With Block Present Almost Continuously for Twenty five Years *Circulation* 1 936 1950
- 230 SCOTT R W and SANCETTA S U Dissecting Aneurysm of Aorta With Hemorrhagic Infarction of the Spinal Cord and Complete Paraplegia *Am Heart J* 38 747 1949
- 231 SØGER M LEQUIME J and DESOLIN H Les effets de l'injection intraveineuse d'acetylcholine chez l'homme L'arrêt des crises de tachycardie paroxystique *Acta Med Scandinav* 122 281 1945
- 232 SIEGAL S and HORN H Quinidine Allergy *Am Heart J* 39 302 1950
- 233 SIGLER L H Hyperactive Cardoinhibitory Carotid Sinus Reflex A Possible Aid in the Diagnosis of Coronary Disease *Arch Int Med* 67 177 1941
- 234 SIMON A J DOLGIN M SOLWAY A J I HIRSCHMANN J and KATZ L N A Re-evaluation of Papaverine in the Treatment of Angina Pectoris *J Lab and Clin Med* 34 992 1949
- 235 SOKOLOW M and EDGAR A L Blood Quinidine Concentrations as a Guide in the Treatment of Cardiac Arrhythmias *Circulation* 1 576 1950
- 236 SORINSON C W and WRIGHT I S A Synthetic Anticoagulant A Polysulfuric Acid Ester of Polyanhydromannuronic Acid (Paritol) Experience with its Use in Man *Circulation* 2 658 1950
- 237 SPINCARN C L and HITZIG W M Orthostatic Circulatory Insufficiency Its Occurrence in Tabes Dorsalis and Addison's Disease *Arch Int Med* 69 23 1942
- 238 SPRAGUE H B and WHITE P D Auricular Flutter Report of a Case of Five Years Duration with Spontaneous Restoration of Normal Rhythm *JAMA* 90 1772 1928
- 239 STARLING L H Some Points in the Pathology of Heart Disease *Lancet* 1 569 1897

- 224 SCHERF D ROMANO F J and TERRANOVA R Experimental Studies on Auricular Flutter and Auricular Fibrillation *Am Heart J* 36 241 1948
- 225 SCHMITZER M A and LEVINE S A Presence of Digitalis in Body Fluids of Digitalized Patients *Arch Int Med* 60 240 1937
- 226 SCHWARTZ S P and BILOON S Clinical Signs of Occluding Thrombi of Left Auricle *Am Heart J* 7 84 1931
- 227 SCHWARTZ S P and DESOLA POOL N Transient Ventricular Fibrillation *Am Heart J* 39 361 1950
- 228 SCHWARTZ S P and LEVI J Digitalis Studies on Children with Heart Disease Effects of Digitalis on Edema of Children With Rheumatic Fever and Chronic Valvular Heart Disease in Presence of Sinus Rhythm *Am J Dis Child* 42 1349 1931
- 229 SCHWARTZ W B and LEVINE S A Case of Paroxysmal Auricular Tachycardia With Block Present Almost Continuously for Twenty five Years *Circulation* 1 936 1950
- 230 SCOTT R W and SANCETTA S U Dissecting Aneurysm of Aorta With Hemorrhagic Infarction of the Spinal Cord and Complete Paraplegia *Am Heart J* 38 747 1949
- 231 SEGER M I EQLINE J and DRNOLIN H Les effets de l'injection intraveineuse d'acetylcholine chez l'homme L'arret des crises de tachycardie paroxystique *Acta Med Scandinav* 122 281 1945
- 232 SIEGAL S and HORN H Quinidine Allergy *Am Heart J* 39 302 1950
- 233 SIGLER I H Hyperactive Cardioinhibitory Carotid Sinus Reflex A Possible Aid in the Diagnosis of Coronary Disease *Arch Int Med* 67 177 1941
- 234 SIMON A J DOLGIN M SOLWAY A J L HIRSCHMANN J and KATZ L N A Re evaluation of Papaverine in the Treatment of Angina Pectoris *J Lab and Clin Med* 34 992 1949
- 235 SOKOLOW M and EDGAR A L Blood Quinidine Concentrations as a Guide in the Treatment of Cardiac Arrhythmias *Circulation* 1 576 1950
- 236 SORFANON C W and WRIGHT I S A Synthetic Anticoagulant A Polysulfuric Acid Ester of Polyanhydromannuronic Acid (Fantol) Experience with its Use in Man *Circulation* 2 658 1950
- 237 STERNBERG C L and HITZIG W M Orthostatic Circulatory Insufficiency Its Occurrence in Tabes Dorsalis and Addison's Disease *Arch Int Med* 60 23 1942
- 238 SPRAGUE H B and WHITE P D Auricular Flutter Report of a Case of Five Years Duration with Spontaneous Restoration of Normal Rhythm *JAMA* 90 1772 1928
- 239 STARLING E H Some Points in the Pathology of Heart Disease *Lancet* 1 569 1897

- 272 WOLFF H G Life Stress and Cardiovascular Disorders *Circulation* 1 187 1950
- 273 WOLFF W I Cardiac Resuscitation Complete Recovery After Over Six Minutes of True Circulatory Arrest *JAMA* 144 738 1950
- 274 WRIGHT I S MARPLE C D and BECK D F Antithrombotic Therapy of Coronary Thrombosis With Myocardial Infarction *JAMA* 138 1074 1948
- 275 WUNSCH R E WARNEKE R D MYERS G B The Effects of Dibenamine on Severe Hypertension *Ann Int Med* 33 613 1950
- 276 WICKOFF J and GOLDRING W Intravenous Injection of Ouabain in Man *Arch Int Med* 39 488 1927
- 277 YOH ~ ~ ~ ~
- 278 YOH ~ ~ ~ ~ osyneph
Tachy
- 279 ZEM ~ ~ ~ ~ Carotid
03 1947
- 280 ZWILLINGER L Über die Magnesia ~ ~ ~ ~ as Herz
Klin Wchnschr 14 1429 1935

Index

- ABDOMINAL belt for postural hypo-
 tension 96
 distention *See* Distention
 pain in digitalis toxicity 21
 in pheochromocytoma 106
 in quinidine therapy 17
 Acetyl choline 32
 Acidosis, in congestive failure 55
 ACTH in coronary occlusion 76
 in rheumatic fever 99
 potassium loss 100
 sodium retention 100
 Addison's disease postural hypo-
 tension in 96
 Adrenal *n. See* Epinephrine
 Agranulocytosis with Pronestyl 37
 Alcohol in angina pectoris 63
 in auricular tachycardia 13 28
 in salt poor diet 53
 paravertebral block 65
 Allergy 32
 danger of Mecholyl 32
 in auricular tachycardia 28
 Alpha Tocopherol (Vitamin E)
 in angina pectoris 64
 Ambulation in myocardial infarc-
 tion 89
 in surgical emergencies 118
 Aminophyllin in Cheyne Stokes
 respiration 51
 in congestive failure 48 55 60
 in coronary insufficiency 67
 in coronary occlusion 74 77 80
 in emergency kit 9
 in nocturnal dyspnea 49
 in pulmonary edema 51
 Ammonium chloride in congestive
 failure 55 60
 Amyl nitrite 96
 danger of syncope 96
 for hiccup 60
 Anemia heart failure in 46 47
 Anesthesia cyclopropane 39 118
 ether 60 118
 in coronary insufficiency 70
 in surgical emergencies 118
 oxygen 118
 pentothal 118
 spinal 118
 Aneurysm arteriovenous 46
 dissecting 108 110
 Angina pectoris 12 61-65
 alcohol 63
 Alpha Tocopherol (Vitamin
 E) 64
 antithyroid drugs 63
 anxiety 67
 biliary colic 117
 cold 62
 coronary insufficiency 62
 differential diagnosis 61
 effort 62
 excitement 62
 in hyperthyroidism 63 64
 in surgical emergencies 117
 I, 63 64
 Kheilin 64
 myocardial necrosis 62
 nitrites 61
 n-troglycerin 62 63 65
 paravertebral block 65
 preceding coronary occlusion
 63
 sedatives 65
 status anginosus 63
 stellate block 65
 subendocardial ischemia 62
 surgical procedures for anasto-
 mosis of aorta to coro-
 nary sinus 65
 omentopexy 65
 thoracic sympathectomy
 65

- Angina pectoris surgical procedures thyroidectomy 63
whiskey 63
- Anticoagulant therapy 80 89
clotting time 84
Depo heparin 84
Dicumarol 84
danger in dissecting aneurysm 108 110
danger in traumatic heart disease 114
heparin 84
paritol 88
prothrombin time 86
thromboplastin 87
transfusion 86
Tromexin 88
urinalysis 86
vitamin K 85 86
- Antidiuretic drug 57
- Antidiuretic hormone in heart failure 45
- Antihistamines in quinidine sensitivity 18
- Antithyroid drugs in angina pectoris 63
- Anxiety in angina pectoris 62
in coronary occlusion 78
in premature beats 44
in syncope 76
- Aortic stenosis calcific 98
syncope in 93
- Aphasia in hypertensive crises 106
- Arrhythmias 11 44
angina in 11
atropin 42
Cedilanid 20
collapse 11
convulsions 39
dibenzamine hydrochloride 39
digitalis 22
digitalis leaf 21
Digitoxin 21 22
Digoxin 20 22
heart failure 11
in coronary occlusion 18 78
in Graves disease 24
in rheumatic fever 99 101
- Arrhythmias in traumatic heart disease 114
magnesium sulfate 39
Ouabain 19
potassium chloride 22 24
Pronestyl 35 79
propyl thiouracil 25
quinidine 24 25
shock 13
strophanthin (K.) 19
syncope 11
vertigo 11
vomiting 11
- Arteriosclerotic heart disease at
nucular fibrillation in 14
- Artificial respiration in surgical emergencies 119, 120
- Ascites in heart failure 48 58 59
- Asthma bronchial coronary insufficiency in 67
danger of Mecholyl 32
- Asthma cardiac 48-49
- Asystole See Cardiac stand still
- Atabrine in ventricular tachycardia 38
- Atelectasis in surgical emergencies 119
- Atropin in A V block 24
in cardiac arrest 119
in emergency kit 9
in pulmonary edema 50
in sinus bradycardia 43 44
in syncope 43 44 93-95
in ventricular fibrillation 42
vs Mecholyl 31
vs parasympathomimetic drugs 37
- Auricular fibrillation 11 12 13
14-25
alcohol 13
A V block 32
chronic 17
digitalis 15 16 27
due to circus movement 14
due to ectopic focus 14
due to Mecholyl 32
embolization 17
emotional disturbance 22

- Auricular fibrillation in arterio-sclerotic heart disease 14
in coronary occlusion 78
in hyperthyroidism 14 15
24 25 32
in pneumonia 14
in rheumatic fever 101
in rheumatic heart disease 15-17 85
in surgical procedures 14 117
118
provysmal 14
potassium salts 19
Pronestyl 34
pulse deficit 14
quinidine 14 16 27
sedation 14
tobacco 13
with heart failure 19
with mitral stenosis 101
- Auricular flutter 11 12 25-28
alcohol 13
A V block 25 32
chronic 26
digitals 26 27
due to ectopic focus 14
esophageal leads 26
in coronary occlusion 78
in surgical emergencies 118
Pronestyl 34
quinidine 27
tobacco 13
venous pulsations in 25
with bundle branch block 25
- Auricular tachycardia 11 12 28-34
alcohol 28
allergy 28
cardiac output 12
carotid sinus 12
differential diagnosis 28
digitals 31
electrocardiogram 12
emotional disturbances 28
gastric distention 28
Graves disease 28
heart failure 12
in surgical emergencies 118
infections 28
- Auricular tachycardia Mechohyl 32
Pronestyl 34
quinidine 31
syncope 12
tobacco 28
- Auricular thrombosis in mitral stenosis 101
- Auricular waves in auricular flutter 25
in esophageal leads 26
- A V block 42, 43, 116
Adrenalin 43
congenital 42
following coronary occlusion 80
digitalis 22 27 42
ephedrine 43
epinephrine 42
idioventricular pacemaker 43
in auricular fibrillation 32
in auricular flutter 25 27 31
in auricular tachycardia 32
in coronary sclerosis 42
Mechohyl 32
puredrine 43
with quinidine 19
shock 42
Stokes Adams syndrome 42
43
syncope 98
- A V conduction effect of carotid sinus pressure 30
effect of eyeball pressure 30
- Azotemia in congestive failure 56
- B
- Bacterial endocarditis subacute 104
- Ball valve thrombus 104
- Basal metabolic rate in paroxysmal tachycardia 12
in pheochromocytoma 106
- Bed rest in coronary occlusion 84
in heart failure 54 58
- Bed tilted in postural hypotension 96

- Benzedrine effect on carotid sinus 30
 Benzodioxane in pheochromocytoma 107
 Beri beri 46 58
 high output failure 46
 Bigeminal rhythm with digitalis 22
 Biliary colic with angina pectoris 117
 Blood volume in cardiac asthma 48
 in cardiac failure 45 47
 hemodilution 48
 sodium retention 47
 Bradycardia *See* Sinus bradycardia
 Bronchospasm due to Mechohyl 32
 in heart failure 48
 Bullet wound 111
 Bundle branch block in auricular flutter 25
 in coronary occlusion 78
 in pulmonary embolism 83
 use of quinidine 19
 in ventricular tachycardia 26
- C
- CALCIUM chloride in cardiac arrest 120
 Cardiac arrest 119 120
 cardiac massage 120
 epinephrine 120
 in surgical emergencies 119 120
 in traumatic heart disease 112
 Cardiac asthma 48-49
 aminophyllin 48
 blood volume 48
 digitalis 49
 diuretics 49
 low sodium diet 49
 morphine 48
 phlebotomy 49
- Cardiac catheterization causing ventricular arrhythmia 34 37
 use of Pronestyl 37
 Cardiac cirrhosis in congestive failure 58 59
 in rheumatic heart disease 58
 Cardiac enlargement in traumatic heart disease 114
 Cardiac failure *See* Heart failure
 Cardiac massage in cardiac arrest 120
 Cardiac output diminished 46
 high 46
 in congestive failure 46 52
 in paroxysmal tachycardia 12
 in postural hypotension 96
 in pulmonary edema 51 52
 in syncope 91 97
 in traumatic heart disease 111
 phlebotomy 46
 salt restriction 46
 Cardiac resuscitation in surgical emergencies 119
 Cardiac standstill 42
 atropine 94
 during Stokes Adams seizures 42
 ephedrine 39 43
 epinephrine 39 43
 in complete A V block 42
 in sinus bradycardia 43
 neosynephrin 37
 paredrine 43
 syncope 93
 in ventricular fibrillation 39
 Cardiac tamponade in rheumatic pericardial effusion 101
 in traumatic heart disease 112
 Carotid sinus benzedrine 30
 convulsions 30
 epinephrine 30
 excision 93
 hemiplegia 30
 in cerebral arteriosclerosis 93
 in supraventricular tachycardia 12 30
 lanatoside C 30
 Mechohyl 30 31

- Carotid sinus neosynephrin 30
 quinine 30
 stimulation 93
 syncope 30 93-95
 tight collars 95
- C
- C
- for congestive failure 57
 hypopotassemia 58
 sodium depletion 58
- Ced land effect on carotid sinus 30
 in arrhythmias 20
 in emergency kit 10
 in pulmonary edema 50
 in supraventricular tachycardia 31
 intravenous 20
- Cerebral anoxia in paroxysmal tachycardia 11
 syncope 91, 97
- Cerebral arteriosclerosis 93
- Cerebral embolism absence of
 hemorrhagic fluid 89
 in coronary occlusion 89
 in rheumatic heart disease 103
- Cerebral symptoms anoxia 91
 in digitalis toxicity 22
- Cervical veins distention in heart failure 48
 pulsations in auricular flutter 25
- Chest injuries non penetrating
 blows 111
 traumatic heart disease 113
- Cheyne Stokes respiration aminophyllin in 51
- Chlorides blood 56
 in salt depletion 56
- Cinchonism 17
- Circus movement in auricular fibrillation and flutter 14
- Clotting time with heparin 84
- CO₂ for hiccup in congestive failure 60
 in surgical emergencies 121
- Codeine 73
- Coffee 13 53
- Cold angina pectoris 62
 coronary insufficiency 69
- Colitis with mercurials 59
- 59
 in coronary occlusion 73
- Contusion of myocardium 114
- Convulsions in arrhythmias 39
 in carotid sinus pressure 30 93
 in hypertensive encephalopathy 106
 in paroxysmal tachycardia 11
 in syncope 97
 in ventricular fibrillation 39
- Coronary artery disease in anesthesia 118
 A V block in 42
 diabetes 70
 insulin in 70
 transfusion in 69
 ventricular tachycardia in 34
 with coronary insufficiency 70
- Coronary insufficiency 12 65-70, 109 113 116
 aminophyllin 67
 anesthesia 70
 angina pectoris 61 62
 bronchial asthma 67
 cold 69
 dehydration 69
 Demerol 67
 diabetes 70
 diarrhea 69
 Dicumarol in 67
 differential diagnosis 65
 digitalis 70
 diuretics 70
 electrocardiogram 65 69 83
 emotional disturbance 69
 fever 66
 heart failure 70
 hemorrhage 70

- Coronary insufficiency hypoglycemia 70
 in traumatic heart disease 114
 leucocytosis 66 67 69
 pulmonary embolism 83
 sedimentation rate 66
 sexual intercourse 69
 shock 69 70
 subendocardial ischemia 66 67
 subendocardial necrosis 69
 surgical procedures 69
 tachycardia 69
 tobacco 69
 transfusions 69 70
 vs coronary occlusion 69 73
- Coronary occlusion 12 44 62 63
 70-90, 85 108 109 113
 ACTH 76
 ambulation 89
 aminophyllin 74 77 89
 anesthesia 90
 anticoagulant therapy 80 84
 arrhythmias 70 78
 arterial embolism 83
 A V block 80
 bundle branch block 78
 caffeine sodium benzoate 89
 codeine 73
 commode 88
 congestive failure 74 77
 cortisone 76
 cough 83
 Demerol 73
 Depo heparin 84
 Dicumarol 84 88 90
 differential diagnosis 70
 digitalis 77
 digoxin 79
 dilaudid 73 76
 distention 73
 duodenal ulcer 79
 electrocardiogram 70 72 78 79
 embolization 73 80 88 89
 embryocardia 70
 enema 88
 epinephrine 76
 fever 72 83
- Coronary occlusion friction rub 73
 gallop rhythm 70
 heparin 84 85 87
 hypertonic glucose 76
 laxatives 88
 leucocytosis 83
 magnesium sulfate 79
 mercurial diuretic 77
 Metapon 74
 morphine 73 76 77 79
 mural thrombi 73
 nausea 70 73
 neosynephrine 76
 danger of nitroglycerin 74
 norepinephrine 76
 obstipation 73
 Ouabain 77
 oxygen 74 76 77 89
 pantopon 74 76
 papaverine 89
 puredrine 76
 paritol 88
 pericarditis 72 73
 perspiration 70
 plasma 76
 premature beats 18 77 78 80
 premonitory phase 72 73
 Priscoline 89
 Pronestyl 78 79
 prothrombin time 84 87
 pulmonary edema 49 70 77
 pulmonary embolism and infarction 80 83
 quinidine 78 79
 salt restriction 77
 sedimentation rate 72 89
 shock 70 78
 sodium amytal 74
 Strophanthin A Strophanthin G 77
 subintimal hemorrhage 72
 surgical emergencies 117
 sympathetic nerve block 89
 syncope 98
 transfusion 76 86
 traumatic 113
 Trendelenburg position 76
 Tromexin 88

- Coronary occlusion urinalysis 86
 ventricular fibrillation 77
 ventricular tachycardia 77
 78 79
 vomiting 70 73
 as coronary insufficiency 65
 69 73
 as dissecting aneurysm 108
- Coronary sinus anastomosis to aorta 65
- Cor pulmonale acute See Pulmonary embolism
- Cor pulmonale chronic cardiac failure 46 47
 high cardiac output 46
- Cortisone in coronary occlusion 76
 in rheumatic fever 99 100
- Cough in coronary occlusion 83
 in pulmonary embolism 83
- Cyanosis in pulmonary edema 49
- Cyclopropane 39, 118
- D**
- DEFIBRILLATION in cardiac arrest 120
- Dehydration in congestive failure 56
 coronary insufficiency in 69
- Demerol antidiuretic effect 57
 coronary insufficiency 67
 coronary occlusion 73
 in emergency kit 10
- Desoxycorticosterone in heart failure 45
 in postural hypotension 96
- Dextrose hypertonic See Glucose
- Diabetes mellitus coronary disease 70
 coronary insufficiency 70
 postural hypotension 96
- Diarrhea coronary insufficiency in 69
 with digitalis 22
 with Mechohyl 31
 with quinidine 17
- Dibenzamine hydrochloride 42 90
 in hypertensive encephalopathy 106
 in paroxysmal tachycardia 39
 in pheochromocytoma 107
 in surgical emergencies 118
 in ventricular fibrillation 42
- Dicumarol 52 84-88, 90
 coronary insufficiency 67
 dangers 85
 hemorrhage 85
 in coronary occlusion 84 88
 in peripheral embolism 103
 in pulmonary infarction 84
 prothrombin time 84 87 103
 urinalysis 86
- Digitalis 19 25
 arrhythmias caused by 22
 A V block caused by 42
 causing supraventricular tachycardia 22
 colored vision 21
 electrocardiogram 22
 in auricular fibrillation 15 17
 27 38
 in auricular flutter 26 27
 in auricular tachycardia 31
 34
 in cardiac asthma 49
 in coronary occlusion 77
 in heart failure 19 46 47 57
 59 70
 in paroxysmal tachycardia 31
 in rheumatic carditis 101
 in rheumatic heart disease 101
 in surgical emergencies 117
 in ventricular tachycardia 38
 intravenous 27 31
 premature beats caused by 34
 toxic effects of 21 24 42 57
 ventricular fibrillation 39
- Digitalis leaf 21
- Digitoxin 21, 22
- Digoxin in arrhythmias 20, 22
 in auricular fibrillation 55
 in congestive failure 52 55
 in coronary occlusion 79

- Digoxin in emergency kit 10
 in pulmonary edema 51
 in rheumatic heart disease 55
 in ventricular tachycardia 37
- Dilaudid for coronary occlusion 73 76
 in emergency kit 10
 in pulmonary edema 50
- Dissecting Aneurysm 108 110
 aortic diastolic murmur 109
 danger of anticoagulants 110
 differential diagnosis 110
 electrocardiogram 109
 hemopericardium 109
 hypertension 109
 mediastinum widened 109
 morphine 110
 occlusion 108
 paraplegia 109
 pericardial effusion 109
 pleural effusion 109
 shock 108 109
 syncope 108
- Distention in auricular tachycardia 28
 in coronary occlusion 73
 in digitalis toxicity 21
 in heart failure 59
 with morphine 73
- Diuretics 54 56
 azotemia 56
 blood chlorides 56
 blood sodium 56
 blood urea 56
 digitalis toxic effects of 57
 in cardiac asthma 49
 in coronary occlusion 77
 in heart failure 47 54 56 57 70 101
 in pulmonary edema 51
 Mercurhydrin 10 51 54 56
 Mercurhydrin ascorbic acid 55
 Mercupurin 51
 Mercuzanthin 55
 muscle cramps 55
 oral 55
 prostate effect on 57
 retention acute 57
- Diuretics salt depletion 56
 salt restriction 57
 Salyrgan 55
 sodium level 56
 Thimerin 10 54 56
 uremia 57
- Dramamine in congestive failure 59
- Dyspnea in congestive failure 48 103
 in pulmonary edema 49
 praxysmal nocturnal 48
 with pulmonary embolism 80
- F
- ECCHYMOSES See Purpura
- Edema 54
 in heart failure 45 48
 pulmonary 49 52
- Effort angina pectoris 62
 pulmonary edema 49
- Elastic bandages in postural hypotension 96
- Electrocardiogram in acute cor pulmonale (pulmonary embolism) 80 83
 in coronary insufficiency 65 69 83
 in coronary occlusion 70 72 78 79
 in dissecting aneurysm 109
 in rheumatic fever 101
 in traumatic heart disease 111 113 114
 potassium 24
- Electroencephalogram in syncope 96
- Electrolytes blood determination 58
 disturbances in congestive failure 57
- Embolism arterial aminophyllin 89
 anticoagulant therapy 80
 caffeine sodium benzoate 89
 cerebral 89
 dibenamine 90

- Embolism arterial embolectomy 90
 femoral pulse 90
 gangrene 90
 in auricular fibrillation, 17
 methonium derivatives 89
 papaverine 89
 peripheral 73 80 89
 Priscoline 89
 saddle embolus 90
 688\ 90
 sympathetic nerve block 89
- Embolism pulmonary 80 83
- Embryocardia in coronary occlusion 70
- Emergency kit, 9
- Emotional disturbances 13
 in coronary insufficiency 69
 hyperthyroidism 12
 in paroxysmal tachycardia 12 13 28
 in pulmonary edema 49
 in sinus tachycardia 43
 syncope 91
- Enemas in congestive failure 59
 in myocardial infarction 88
- Ephedrine antidiuretic effect 57
 in A V block 43
 in Stokes Adams syndrome 43
 in surgical emergencies 117 118
 in syncope 43 93 95
- Epinephrine antidiuretic effect 57
 effect on carotid sinus 30
 in A V block 42
 in coronary occlusion 76
 in emergency kit 10
 in pheochromocytoma 106
 in Stokes Adams syndrome 43
 in surgical emergencies 119 120
 syncope 91
- Esophageal leads in auricular flutter 26
- Esophageal spasm as angina pectoris 61
- Etamon See Tetra ethyl ammonium chloride
- Ether for hiccup 60
- Fthal chloride for hiccup 60
- Eye-ball pressure effect in A V conduction 30
 in supraventricular tachycardia 30
- F
- FAINTING See Syncope
- Fever in coronary insufficiency 60
 in coronary occlusion 72
 with pulmonary embolism 80
- Fluids intravenous in surgical emergencies 118
 restriction of 59
 retention of 45
- Flutter waves 25
- Foci of infection in surgical emergencies 117
- Foods restriction in congestive failure 59
- Friction rub in coronary occlusion 73
 in rheumatic fever 101
- Fruit juices 53
 cause of nausea in congestive failure 60
 in rheumatic fever 100
- G
- GALLBLADDER disease effect of operation on angina 117
 as angina pectoris 61
- Gallop rhythm in coronary occlusion 70
- Gangrene ice packs 90
- Ginger ale 53
 for nausea in congestive failure 50
- Glucose hypertonic, in coronary occlusion 76
 in hypertensive encephalopathy 106
 in syncope 97
 in traumatic heart disease 112

- Glucose tolerance test in syncope 97
- Glucosuria in pheochromocytoma 106 107
- Grand mal syncope 91
- Graves disease *See* Hyperthyroidism
- H**
- HEADACHE 17
- in angina pectoris 62
- in hypertension 106
- Heart block *See* A V block
- Heart failure 45-61, 101 116
- abdominal distention 59
- acidosis 55
- aminophyllin 48 55 60
- ammonium chloride 55 60
- anemia 46 47 58
- apnoea of bone marrow 45
- antidiuretic hormone 45
- drugs 47
- arrhythmias 11 13
- ascites 48 58 59
- auricular fibrillation 14 19 101
- azotemia 56
- bed rest 54 58
- beri beri 58
- blood electrolytes 57
- urea 56
- volume 45 47 48
- bloodless phlebotomy 49
- bronchospasm 48
- cardiac asthma 48
- arrhythmias 58 59
- output 46 47 51
- cervical veins distention of 48
- chloride level 56
- colitis 59
- cation exchange resins for 57
- constipation 59
- cor pulmonale 47
- coronary occlusion 76
- dehydration 56
- Demerol 57
- desoxycorticosterone 45
- Heart failure Diurematol 52
- digitalis 19 31 46 49 57 59 70
- Digoxin 52 55
- diuretics 56 57
- Dramamine 59
- dyspnea 48
- edema 45 48
- enemas 59
- fluid intake 53
- hemodilution 48
- hepatomegaly 48
- hiccup 59 60
- hydrothorax 58
- hypertensive heart disease 47
- hyperthyroidism 17 58
- hypopotassemia 55
- in rheumatic fever 90 100 101
- in rheumatic heart disease 53 58 103
- in sinus tachycardia 43
- in traumatic heart disease 111 114
- linoleic 53
- laxative 54 59
- left ventricular failure 49 52 79
- liver disease 59
- mechanism (pathogenesis) 45
- backward failure 45 46
- cardiac output 46 47 51
- forward failure 45 46
- Mercuhydram 54 56
- mercurial diuretic 47 49 55 101
- Mercuzanthin 55
- morphine 48 49 57 60
- nausea 57 59
- obesity 54
- opiates 59
- orthopnea 48
- Pantopon 50
- paracentesis 58 59
- penicillin 52
- phlebotomy 46 47
- potassium 57
- protein intake 59
- pulmonary congestion 48

- Heart failure pulmonary disease*
 chronic 47
 infarction 52
 infection 52
 renal blood flow 47 48
 salt depletion 56 58
 poor diet 54 58
 restriction 46 53 57 101
 substitutes 53
 shock 103
 sodium depletion 49 52
 excretion 53
 level 56
 retention 47 48 54
 stroke volume 46
 surgical emergency
 sympathetic 49
 tachycardia
 therapy 52 60
 Thrombin 54 56
 thoracentesis 58
 tourniquets 49
 uremia 57
 vitamins 59
 vomiting 57 59
 weight reduction 54
 wheezing 48
Heart wound 111
Hematuria due to quinidine 18
Hemiplegia in carotid sinus pressure 30
Hemoglobin 52
Hemopericardium in dissecting aneurysm 109
 in traumatic heart disease 111 112 115
Hemoptysis in mitral stenosis 105
 in pulmonary embolism 83
Hemorrhage as surgical emergency 117 121
 subintimal in coronary occlusion 72
 syncope in 91
 transfusion 69
Heparin 88
 clotting time 84 87
 coronary occlusion 85 87
 in pulmonary infarction 84
Hepatic disease in congestive failure 59
 syncope in 97
Hepatomegaly in heart failure 48
Herpes zoster in angina pectoris 61
Hiatus hernia in angina pectoris 61
Hiccup in heart failure 59 60
 treatment of 60
Histamine in pheochromocytoma 107
Homan's sign in venous thrombosis 83
Hunger in venous thrombosis 83
 syncope 97
Hyperpnea 56
Hypertension 106 114
 and dissecting aneurysm 108 109
 encephalopathy 106
 in pheochromocytoma 107
 malignant 106
Hypertensive encephalopathy 106 107
 aphasia 106
 convulsions 106
 dextrose 106
 Dibenzamine 106
 headache 106
 in pheochromocytoma 106
 magnesium sulfate 106
 nephritis acute 106
 paralysis 106
 spinal tap 106
Hypertensive heart disease 46
 anesthesia in 118
 cardiac failure in 47
Hyperthyroidism angina pectoris 63 64
 auricular fibrillation 14 24, 25
 congestive failure 47 58
 digitalis 22 24
 high cardiac output 46

- Hyperthyroidism Lugol's solution 24
 Methylol 32
 propylthiouracil 25
 quinidine 24 25
 radioactive iodine I_{131} 22 24 63 64
 supraventricular tachycardia 12 28
- Hyperventilation syncope 96
- Hypoglycemia in coronary insufficiency 70
 in syncope 97 98
- Hypopotassemia 57 58
 cation exchange resins 58
 in congestive failure 55
- Hypotension See Postural hypotension 95
- Hypothyroidism induced in angina pectoris 63
- Hysteria syncope 96
- I**
- I_{131} See Radioactive iodine
- Ice packs gangrene 90
- Idioventricular pacemaker Stokes Adams syndrome 42
- Infarction in traumatic heart disease 112
 myocardial 62
 pulmonary 83 118 119
 in angina pectoris 62
- Infections supraventricular tachycardia 28
- Intratracheal tube in cardiac arrest 120
- Ipecac syrup of in emergency kit 10
 in supraventricular tachycardia 31
- Ischemia myocardial in coronary insufficiency 66
- J**
- Joints effusions in rheumatic fever 100
- Joints pains in rheumatic fever 100
 silylates 100
- K**
- KERLIN angina pectoris 64
 nausea and vomiting 64
- L**
- LAVATOSIDE C See Cedi and Lanolac powder 53
- Laxatives cascara sagrada 59 88
 in heart failure 54 59
 in myocardial infarction 88
 milk of magnesia 59 88
 mineral oil 88
- Lethargy in salt depletion 56
- Leucocytosis in coronary insufficiency 66 67 69
 in pulmonary infarction 83
- Liver disease in congestive failure 59
- Lugol's solution 24 25 64
- M**
- MAGNESIUM sulfate in coronary occlusion 79
 in emergency kit 10
 in hypertensive encephalopathy 106
 in ventricular fibrillation 39
 in ventricular tachycardia 39
 intravenous 38
 nausea 38
 vertigo 38
 weakness 38
- Meds in arrhythmias 13
 in pulmonary edema 49
 small 53
- Methylol 30 32
 allergy 32
 asthma 32
 atropin 31

- Mecholyl auricular tachycardia 32/
 reactions to 32
 Mediastinum widened in dissect-
 ing aneurysm 109
 Mercurhydrin *See* Diuretics
 Mercurhydrin ascorbic acid 54 79
 Mercupurin *See* Diuretics
 Mercurial diuretics *See* Diuretics
 Mercuzanthin *See* Diuretics
 Metapon in coronary occlusion 74
 Methonium derivatives (C_3 , C_4) 89
 Milk nausea due to 60
 Milk of magnesia in congestive
 failure 59
 in myocardial infarction 88
 Mineral oil in myocardial infarc-
 tion 88
 Mitral stenosis 16 101-105
 anticoagulant therapy 103
 auricular fibrillation 101
 auricular thrombi 101
 ball valve thrombus 103
 hemoptysis 105
 left auricular appendectomy
 103
 operations on mitral valve
 105
 peripheral embolism 103
 pulmonary edema 105
 sympathetic block 105
 tight 104
 Morphine antidiuretic effect 57
 in coronary occlusion 74 76
 77 79
 in dissecting aneurysm 110
 in emergency kit 10
 in heart failure 48 49 60 103
 in acute pericarditis 101
 in pulmonary edema 49 50
 in traumatic heart disease 112
 in ventricular tachycardia 38
 intravenous 38
 Myocardial infarction *See* Coro-
 nary occlusion
 Myocardium 111 112
 contusion 114
 infarction *See* Coronary oc-
 clusion
 laceration 111 112
- N
- Nausea congestive failure 59
 coronary occlusion 70 73
 dig talis 22 57
 Khellin 64
 magnesium sulfate 38
 Mecholyl 31
 morphine 73
 pheochromocytoma 106
 Pronestyl 37
 pyridoxine for 59
 quinidine 17
 syncope 91
 treatment 59
 Neostigmine 32
 Neosynephrin cardiac standstill 37
 carotid sinus 30
 coronary occlusion 76
 danger in ventricular prema-
 ture beats 34
 emergency kit 10
 nausea 37
 supraventricular tachycardia
 32
 surgical emergencies 117 118
 syncope 95
 Nephritis acute 106
 Neurocirculatory asthenia postu-
 ral hypotension 95
 Niacin intravenous for hiccup
 in congestive failure 60
 Nitrites 61
 Nitroglycerin 96
 in angina pectoris 62 63 64
 in coronary occlusion 74
 in emergency kit 10
 flushes 62
 headache 62
 syncope 62
 throbbing 62
 Nodal tachycardia *See* Auricular
 tachycardia
 Nor epinephrine in coronary oc-
 clusion 76
 in hypertensive encephalop-
 athy 106

O

- OBESITY in heart failure 54
in surgical emergencies 119
- Occlusion arterial cerebral 103
chronic dicumarolization 103
in dissecting aneurysm 108
in rheumatic heart disease 103
- Operative procedures *See* Surgical procedures
- Orthopnea in heart failure 48
- Oubain *See* Strophanthin G
- Oxygen therapy in congestive failure 103
in coronary occlusion 74 76 77 89
in hiccup with congestive failure 60
in pulmonary edema 50
in surgical emergencies 118 121

P

- PACLET'S disease 46
- Palpitation 12
in pheochromocytoma 106 107
- Pancarditis 99
- Pantopon coronary occlusion 74 76
pulmonary edema 50
- Papaverine in angina pectoris 64
in embolization 89
intraarterial 89
- Paracentesis abdominis in congestive failure 58 59
- Paralysis hypertensive crises 106
- Paraplegia in dissecting aneurysm 109
- Parasympathomimetic drugs *see* cholinergics
atropine with 32
in supraventricular tachycardia 32
- Mecholyl 30 32
neostigmine 32

- Paravertebral block alcohol 65
in angina pectoris 65
in mitral stenosis 105
tetra ethyl ammonium chloride 65
- Paredrine in coronary occlusion 76
in Stokes Adams syndrome 43
in syncope 95
- Paristol 88
- Paroxysmal ventricular fibrillation *See* Atrial fibrillation
- Paroxysmal tachycardia *See* Ventricular fibrillation Atrial flutter Atrial tachycardia
- Penicillin in heart failure 52
- Pentothal 118
- Pepper 53
- Peptic ulcer in angina pectoris 61
- Pericardial effusion in dissecting aneurysm 109
in pericardial tip 111
in rheumatic fever 100
in traumatic heart disease 113
subacute 100
- Pericardial rub in traumatic heart disease 112 114
- Pericarditis acute 99
in rheumatic fever 99 101
in traumatic heart disease 113
in coronary occlusion 73
- Peripheral resistance syncope 91
- Pernicious anemia heart failure in 58
- Perspiration *See* Sweating
- Petechiae white centered in bacterial endocarditis subacute 104
- Pheochromocytoma 106 107
abdominal pain 106
basal metabolic rate 106
Benzodioxane 107
Dibenzamine 107
epinephrine 106 107
glycosuria 106 107
histamine 107
hypertensive encephalopathy 106
nausea 106

- Pheo chromocytoma nor epineph
 rine 106
 palpitation 106 107
 postural hypotension 96
 sweating 106 107
 tachycardia 106
 vertigo 106
 vomiting 106
 Phlebotomy bloodless (tourni-
 quets) 49 50 51
 in heart failure 16 47
 in pulmonary edema 51 52
 needles for in emergency kit
 10
 Plasma in coronary occlusion 76
 in traumatic heart disease 112
 Pleural effusion in dissecting
 aneurysm 109
 in heart failure 58
 Pleurisy vs angina pectoris 61
 Pneumonia auricular fibrillation
 in 14
 Pneumothorax vs angina pectoris
 61
 Polomyelitis pulmonary edema
 in 50
 Postural hypotension 95 96
 Addison's disease 96
 amyl nitrite 96
 cardiac output 96
 cord tumor 95
 diabetes mellitus 96
 hysteria 96
 neurocirculatory asthenia 96
 nitroglycerin 96
 pheochromocytoma 96
 syncope 93 96
 tabes dorsalis 95
 treatment abdominal belt 96
 atropin 96
 bed tilted 96
 desoxycorticosterone 96
 elastic bandages 96
 epinephrine 96
 neosynephrin 96
 paredrine 96
 salt 96
 varicose veins 96
 Potassium blood determination
 100
 effect on electrical discharges
 24
 in congestive failure 1
 loss with ACTH 100
 Potassium salts fruit juice 100
 in auricular fibrillation 10, 22
 in emergency kit 10
 in ventricular tachycardia 38
 toxic effects of 24
 P R interval prolonged coronary
 occlusion 80
 digitalis 22
 Premature beats 43 44, 80 114
 anxiety 44
 coronary occlusion 44 77 78
 digitalis 34
 in congestive failure 57
 in mitral stenosis 16
 in rheumatic heart disease 16
 in surgical procedures 44
 neosynephrine 34
 paroxysmal tachycardia 44
 Pronestyl 44
 psychiatric therapy 44
 ventricular 18 22 34 35
 Prisolone in peripheral embolism
 intraarterial 89
 Procaine intravenous during sur-
 gical procedures 37 119
 120
 Procaine amide see Pronestyl
 Pronestyl (procaine amide) 35 38
 agranulocytosis 37
 auricular fibrillation 34
 flutter 34
 tachycardia 34
 cardiac catheterization 37
 standstill 38
 coronary occlusion 78 80
 emergency kit 10
 intravenous 42
 prophylactic 38
 purpura 37
 surgical procedures 37 118
 ventricular fibrillation 42
 premature beats 44
 tachycardia 35 38

- Propyl thiouracil in arrhythmias 25
 Prostatic disease effect of mercurials 57
 Protamine sulfate in heparinization 85
 Protein intake in congestive failure 59
 Prothrombin time 84-87, 103
 bedside method 87
 in Dicumarol therapy 84 87 103
 Psychiatric therapy premature beats 44
 syncope 96
 Pulmonary congestion amino-phyllin 77
 in coronary occlusion 70
 in heart failure 48
 mercurial diuretic 77
 morphine 77
 oxygen 77
 Pulmonary disease chronic cardiac failure 46 47
 Pulmonary edema acute 49 52
 aminophyllin 50
 atropin 50
 bubbling rules 49
 cardiac output 51 52
 Cheyne Stokes respiration 51
 cyanosis 49
 Digoxin 51
 Dilaudid 50
 diuretics mercurial 51
 dyspnea 49
 emotional disturbances 49
 exercise 49
 hematocrit 52
 in coronary occlusion 49 70 77
 in mitral stenosis 105
 in surgical emergencies 118
 meals large 49
 morphine 49 50 77
 Ouabain 51
 oxygen 50 77
 packed red blood cells 52
 Pantopon 50
 Pulmonary edema paroxysmal tachycardia 49
 phlebotomy 50
 bloodless 49 51 52
 poliomyelitis 50
 shock 49 50
 strophanthin (K. & G.) 36 51
 sympathetic block 105
 triuran 49
 venous pressure 52
 wheezing 50
 Pulmonary embolism 80-83, 116
 anticoagulant therapy reduced by 80
 right bundle branch block 83
 coronary insufficiency 83
 acute cor pulmonale 81
 cough 83
 dyspnea 80
 electrocardiogram 80
 fever 80
 hemoptysis 83
 Homans sign 83
 shock 80
 tachycardia 80
 Pulmonary infarction anticoagulant therapy 52 84
 dissecting aneurysm 107
 fever 83
 in congestive failure 52
 in rheumatic heart disease 103
 leucocytosis 83
 Pulmonary infection in congestive failure 52
 penicillin 52
 Pulse deficit in auricular fibrillation 14
 Purpura effect of quinidine 17 18
 in rheumatic heart disease 85
 Pronestyl 37
 Pyridoxine in nurse in congestive failure 59

Q

QUINIDINE 14-19, 30 31

- Quinidine antidiuretic effect 57
 antihistamines 18
 carotid sinus 30
 cinchonism 17
 encephalopathy of 16
 in auricular fibrillation 14 16
 in auricular flutter 27
 in auricular tachycardia 31
 in A V block 19
 in bundle branch block 19 78
 in coronary occlusion 78 79
 in emergency kit 10
 in hiccup 60
 in premature beats 18 44 80
 in supraventricular tachycardia 34
 in surgical emergencies 118
 in ventricular fibrillation 38 39
 in ventricular tachycardia 35
 idiosyncrasy 17
 intramuscular 15 27 31 78
 intravenous 15 39
 maintenance dose 15 16 27
 Mechohyl 32
 plasma concentration 14 15
 prophylactic 16 38
 sensitivity 14 17 18 38
 toxic effects 17
 electrocardiographic changes 18
 hematuria 18
 laryngeal edema 18
 purpura 18
 Quinine in ventricular tachycardia 79
- R
- Radioactive iodine in angina pectoris 63 64
 in hyperthyroidism 22 24, 25,
 Renal function blood flow 45 47 48
 excretion 47
 Respiration artificial during operation 119, 120
 Cheyne Stokes as nophyllin 51
 depressed by quinidine 17
- Rheumatic heart disease 14-16 34
 auricular fibrillation 15 50 85 101
 bacterial endocarditis subacute 104
 ball valve thrombus 103
 blood culture 104
 cardiac cirrhosis 58
 congestive failure 58
 digitalis 17
 Digoxin 55
 dyspnea 103
 hemoptysis 105
 lanatoside C 50
 left auricular appendectomy 103
 mercurial injections 55
 mitral stenosis 16 101 104
 morphine 103
 oxygen 51 103
 petechiae 104
 phlebotomy bloodless 50
 premature beats 16
 pulmonary edema 50
 pulmonary infarction 103
 purpura 85
 shock 103
 splenomegaly 104
 surgical emergencies 105 117
 ventricular tachycardia 44
 Rheumatic fever 99 101
 ACTH 99
 arrhythmias 99 101
 auricular fibrillation 101
 blood potassium 100
 cardiac tamponade 101
 congestive failure 99 101
 cortisone 99 100
 digitalis 101
 electrocardiogram 101
 friction rub 101
 fruit juices 100
 joint effusions 100
 joint pains 100
 morphine 101
 pericarditis 99
 pericardial effusion 100
 pericarditis acute 99 101

- Rheumatic fever salicylates 99
 100
 sedimentation rate 100
 shock 101
 Rupture of heart 113
 of valve 111
- S
- SALICYLATES in rheumatic fever
 99 100
 Saline intravenous in surgical
 emergencies 118
 Salivation with Mecholyl 31
 Salt depletion with diuretics 56
 in heart failure 46 53 54
 in postural hypotension 96
 restriction 46 47 53 54 58
 77
 Salt substitutes in heart failure
 53
 potassium chloride 53
 Salyrgan 55
 Scotomata digitalis therapy 22
 Sedatives hiccup in congestive
 failure 60
 in angina pectoris 65
 in auricular fibrillation 14
 in paroxysmal tachycardia 13
 in syncope 97
 Sedimentation rate in coronary
 insufficiency 66
 in coronary occlusion 72 89
 in rheumatic fever 100
 Septal defect interatrial creation
 in mitral stenosis 103
 Sexual intercourse coronary in
 sufficiency 69
 Shock 31 78
 Adrenalin 42
 epinephrine for 42
 in arrhythmias 13
 in A V block 42
 in congestive failure 103
 in coronary occlusion 70
 in dissecting aneurysm 108
 109
 in paroxysmal tachycardia 12
 Shock in pulmonary edema 49 50
 in pulmonary embolism 80
 in rheumatic fever 101
 in traumatic heart disease
 111 112
 in ventricular fibrillation 39
 Mecholyl 31
 morphine sulfate for 50
 oxygen 103
 peripheral 49
 postoperative in coronary in
 sufficiency 69
 Trendelenburg position 76
 Sinus bradycardia 43
 atropin 43 93
 carotid sinus stimulation 93
 digitalis 22
 dizziness 43
 ephedrine 43 93
 syncope 43
 Sinus tachycardia See Tachy-
 cardia
 688 \ 90
 Skin eruptions with quinidine 17
 Sodium blood level 56
 depletion 58
 diet low in 49 52 100
 excretion 45 47 53
 in congestive failure 47, 49 52 54
 in surgical emergencies 118
 retention 45 47, 48 54 100
 Sodium amytal in coronary occlu-
 sion 74
 in emergency kit 10
 Sodium bicarbonate 54
 Spinal cord tumor postural hypo-
 tension in 95
 Spinal tap in hypertensive enceph-
 alopathy 106
 Splenomegaly in subacute bac-
 terial endocarditis 104
 Spondylitis re angina pectoris 61
 Stab wound of heart 111 112
 Status anginosus 63
 Steering wheel accidents in
 traumatic heart disease 114
 Stellate block in angina pectoris
 with ammonium sulfate 65
 with status anginosus

- Stoke-Adams syndrome ephedrine 43
 epinephrine 43
 idioventricular pacemaker 42
 in heart block 42 43
 in ventricular fibrillation 39
 paredrine 43
 syncope 98
- Stroke volume in cardiac failure 46
- Strophanthin (K & G) auricular fibrillation 19 20
 coronary occlusion 77
 emergency kit 10
 pulmonary edema 36 51
 supraventricular tachycardia 31
- Subendocardium ischemia in angina pectoris 62
 in coronary insufficiency 67
 necrosis 69
- Sudden death in ventricular fibrillation 39
- Supraventricular tachycardia *See* Auricular tachycardia
- Surgical emergencies 116-121
 age 117
 ambulation 118
 anesthesia 118
 arrhythmias 117 118
 artificial respiration 119 120
 cardiac resuscitation 119
 cholecystectomy 117
 CO 121
 coronary occlusion 117
 defibrillation 120
 digitalization 117
 ephedrine 117 118
 epinephrine 119 120
 fluid intake 118
 heart failure 117 118
 hemorrhage 117 121
 intravenous saline 118
 neosynephrine 117 118
 oxygen 118 121
 procaine 119 120
 Pronestyl 118
 pulmonary edema 118
 infarction 118 119
 quinidine 118
- Surgical emergencies rheumatic heart disease 117
 Stokes Adams syndrome 117
 transfusion 117 121
 ventricular fibrillation 120
 with angina pectoris 117
- Surgical procedures anastomosis of aorta to coronary sinus 65
 auricular appendectomy 103
 embolectomy 90
 excision of carotid sinus 95
 on mitral valve 105
 omentopexy 65
 phrenicectomy 60
 stellate ganglion block 65
 thoracic sympathectomy 65
 thyroidectomy 63
 traumatic heart disease 112
- Sweating in coronary occlusion 70
 in pheochromocytoma 106 107
 syncope 91
- Sympathectomy thoracic in angina pectoris 65
- Sympathetic nerve block 89 90
- Syncope 30 32 91-98
 angina pectoris 62
 anxiety 96
 aortic stenosis 93 98
 arrhythmias 11
 asystole 93
 cardiac output 91 97
 carotid sinus 30 93 95
 cerebral anoxia 91
 arteriosclerosis 93
 clonic twitching 91
 convulsions 95 97
 dizziness 97
 electroencephalogram 96
 emotional disturbances 91
 epinephrine 91
 fright 91 93
 glucose tolerance test 97
 grand mal 91
 hemorrhage 91
 hepatic disease 97
 hunger 97
 hyperventilation 96
 hypoglycemia 97 98

- Syncope hysterical 96
 in dissecting aneurysm 108
 neurogenic 91
 overactive pancreas 97
 pancreatic adenomata 97
 peripheral resistance 91
 post coronary occlusion 98
 postural hypotension 93 95
 sight of blood 91 93
sinus bradycardia 43
 Stokes Adams syndrome 43 98
 straining at stool 97
 suggestion 96
 tight collars 95
 treatment atropin 93 95
 ephedrine 93 95
 excision of carotid sinus 95
 glucose 97
 neosynephrine 95
 Paredrine 95
 psychiatric 96
 sedation 97
 Valsalva maneuver 97
 vasodepressor 91 93

T

- TABES dorsalis* postural hypotension 95
 Tamponade cardiac in rheumatic fever 101
 in traumatic heart disease 111 112
 Tetraethyl ammonium chloride (Etamon) 65 89
 in pheochromocytoma 107
 Thiomerin in emergency kit 10
 in heart failure 54 56
 Thoracentesis in congestive failure 58
 Thromboplastin 87
 Thrombosis auricular 101
 ball valve thrombus 103
 coronary See Coronary occlusion
 mural in coronary occlusion 73
 in mitral stenosis 101
 venous Homan's sign in 83
 Thyroidectomy in angina pectoris 63
 Thyrotoxicosis See Hyperthyroidism
 Tinnitus 17
 Tobacco in auricular tachycardia 28
 in coronary insufficiency 69
 supraventricular tachycardia 13 28
 Tourniquets in bloodless phlebotomy 49 51
 in emergency kit 10
 Transfusion 18
 in anticoagulant therapy 86
 in coronary disease 69
 in coronary insufficiency 69 70
 in coronary occlusion 76
 in pulmonary edema 52
 with packed red blood cells 52
 in surgical emergency 117 121
 in traumatic heart disease 112
 intraarterial 121
 Trendelenburg position for shock 76
 in traumatic heart disease 112
 Trigeminal rhythm with digitals 22
 Tromexan 88

U

- URFA blood during mercurial diuretic therapy 56
 Uremia caused by mercurial diuretics 57
 Urinolysis in Dicumarol therapy 86
 Urinary retention caused by mercurials 57

V

- Vagal stimulation breath holding in supraventricular tachycardia 28
 in surgical emergencies 119
 Valsalva maneuver in syncope 97

- Varicose veins, causing postural hypotension 96
- Venous pressure 45
in pulmonary edema 52
- Venous pulsations auricular flutter 25
in ventricular tachycardia 34
- Ventricular fibrillation 11 39-42
atropin 42
convulsions 39
Dibenzamine 42
digitalis 39
in coronary occlusion 39 77
in surgical emergencies 120
magnesium sulfate 39
Pronestyl 42
quinidine 39
shock 39
Stokes Adams syndrome 39
sudden death 39
- Ventricular tachycardia 11 12
34-39, 79
Atabrine 38
A V block 42
coronary disease 13 34
digitalis 22 38
during cardiac catheterization 34
in coronary occlusion 13 77 79
in rheumatic heart disease 34
magnesium sulfate intravenous 38
Mercurhydrin 37
morphine 38
potassium chloride 22
salts 38
Pronestyl 37 38 78
quinidine 35 38 78
venous pulse tracings 34
vs bundle branch block 26
- Vertigo 22
in arrhythmias 11
with magnesium sulfate 38
in pheochromocytoma 106
- Vertigo in syncope 97
- Visual disturbances with digitalis 21
with quinidine 17
- Vitamin L (Alpha Tocopherol) in angina pectoris 64
- Vitamin K in Dicumarol therapy 85 86
- Vitamin K₁ oxide 86
- Vitamins intake in congestive failure 59
- Vomiting 31
digitalis toxicity 57
Dramamine for 59
in arrhythmias 11
in congestive failure 59
in coronary occlusion 70 71
in pheochromocytoma 106
pyridoxine for 59
with digitalis 22
with Kheslin 64
with Mechohyl 31
with morphine 73
with quinidine 17
treatment 59
- W
- WATER excretion 47
intake 53
- Weakness in salt depletion 56
in syncope 97
with magnesium sulfate 38
- Weight loss 54
reduction in heart failure 54
- Wheezing in cardiac asthma 48
in pulmonary edema 50
- Y
- Y RAY in dissecting aneurysm 109
in traumatic heart disease 111